arXiv:1202.0875v1 [math.DG] 4 Feb 2012

SMOOTH CONVERGENCE AWAY FROM SINGULAR SETS

SAJJAD LAKZIAN AND CHRISTINA SORMANI

Abstract. We consider sequences of metrics, g_j, on a Riemannian manifold, M, which converge smoothly on compact sets away from a singular set $S \subset M$, to a metric, g_∞, on $M \setminus S$. We prove theorems which describe when $M_j = (M, g_j)$ converge in the Gromov-Hausdorff sense to the metric completion, (M_∞, d_∞), of $(M \setminus S, g_\infty)$. To obtain these theorems, we study the intrinsic flat limits of the sequences. A new method, we call hemispherical embedding, is applied to obtain explicit estimates on the Gromov-Hausdorff and Intrinsic Flat distances between Riemannian manifolds with diffeomorphic subdomains.

1. Introduction

The purpose of this paper is to provide general results concerning the limits of Riemannian manifolds which converge smoothly away from a singular set as follows:

Definition 1.1. *We will say that a sequence of Riemannian metrics g_j on a compact manifold M converges smoothly away from $S \subset M$ to a Riemannian metric g_∞ on $M \setminus S$ if for every compact set $K \subset M \setminus S$, g_j converge $C^{k,\alpha}$ smoothly to g_∞ as tensors.*

The techniques developed in this paper will also be applied to other notions of smooth convergence away from singular sets in upcoming work of the first author, particular notions in which the sequence of manifolds need not be diffeomorphic. With any notion of smooth convergence away from a singular set, one must keep in mind that even when the singular set is an isolated point, smooth convergence away from that point does not even imply that (M_∞, g_∞) is compact [Example 3.11]. Increasingly large distances may exist outside the compact sets used to define the smooth convergence.

Given two compact Riemannian manifolds, M_i, the Gromov-Hausdorff distance, $d_{GH}(M_1, M_2)$, is an isometry invariant. Introduced by Gromov in [Gro99], it is a distance on compact metric spaces in the sense that $d_{GH}(M_1, M_2) = 0$ iff M_1 is isometric to M_2. When studying precompact domains within manifolds, one always takes the metric completion before examining the region using the Gromov-Hausdorff distance. Section 2 (see Definition 2.5).

Lakzian was partially supported as a doctoral student by NSF DMS #1006059.

Sormani was partially supported by NSF DMS #1006059.

Smooth limits away from singular sets, depend on the charts and tensors g_j used to define the smooth limit (c.f. Example 3.7). Thus it is important to understand when the metric completion, $\bar{Y}$, of a smooth limit, $Y = (M \setminus S, g_\infty)$, is in fact actually the Gromov-Hausdorff limit, (M_0, d_0), of the original sequence of manifolds, (M_j, d_j), where d_j is the Riemannian distance defined by the Riemannian metric g_j.. Observe that these spaces need not be isometric (c.f. Example 3.1) and that the original sequence of manifolds might not even have a Gromov-Hausdorff limit (c.f. Example 3.10). If $M \setminus S$ is not connected there isn't even a notion of the metric completion as a single metric space (c.f. Example 3.4).

Theorems relating Gromov-Hausdorff limits and smooth limits away from singular sets appear in work of Anderson, Bando-Kasue-Nakajima, Eyssidieux-Guedj-Zeriahi, Huang, Ruan-Zhong, Sesum, Tian and Tosatti particularly in the setting of Kahler Einstein manifolds [And89] [BKN89] [EGZ09] [Hua09] [RZ] [Ses04] [Tia90] [Tos09]. However, even in this setting, the relationship is not completely clear and the limits need not agree [Ban90].

In this paper, our primary goal is to examine when the metric completion, (M_∞, d_∞), of the smooth limit, $(M \setminus S, g_\infty)$, is isometric to the Gromov-Hausdorff limit, (M_0, d_0), of the original sequence of Riemannian manifolds (M, g_j). We prove a number of theorems and present a number of examples considering manifolds with and without Ricci curvature bounds. Perhaps the most important result is the following:

Theorem 1.2. *Let $M_i = (M, g_i)$ be a sequence of oriented Riemannian manifolds with uniform lower Ricci curvature bounds,*

$$\mathrm{Ricci}_{\mathrm{g_i}}(\mathrm{V}, \mathrm{V}) \geq (\mathrm{n}-1)\mathrm{H}\, \mathrm{g_i}(\mathrm{V}, \mathrm{V}) \qquad \forall \mathrm{V} \in \mathrm{TM_i} \tag{1}$$

which converges smoothly away from a singular set, S, of codimension two. If there is a connected precompact exhaustion, W_j, of $M \setminus S$,

$$\bar{W}_j \subset W_{j+1} \text{ with } \bigcup_{j=1}^{\infty} W_j = M \setminus S \tag{2}$$

satisfying

$$\mathrm{Vol}_{g_i}(\partial W_j) \leq A_0 \tag{3}$$

and

$$\mathrm{Vol}_{g_i}(M \setminus W_j) \leq V_j \text{ where } \lim_{j\to\infty} V_j = 0 \tag{4}$$

then

$$\lim_{j\to\infty} d_{GH}(M_j, N) = 0 \tag{5}$$

where N is the metric completion of $(M \setminus S, g_\infty)$.

Note that, unlike prior existing results concerning the Gromov-Hausdorff limits of manifolds, here we require only area and volume controls on the connected precompact exhaustion. Theorem 1.2 is a consequence of Theorem 6.10, stated within, which assumes only that the connected precompact exhaustion is uniformly well embedded in the sense of Definition 5.1. The necessity of the various hypothesis of these theorems is described in Remark 6.13.

The Ricci curvature condition in these theorems may be replaced by a requirement that the sequence of manifolds have a uniform linear contractibility function. See Definition 6.1, Theorem 6.7 and Theorem 6.6, stated within. The necessity of the various hypothesis of these theorems is described in Remark 6.8.

Observe our main theorems concern sequences of manifolds converging smoothly away from a singular set satisfying (3) and (4). In order to control the limits of such manifolds using only conditions on volumes, we apply techniques developed by the second author with Stefan Wenger in [SW10] and [SW11]. In attempt to keep this article self contained, we review convergence of Riemannian manifolds in Section 2. We provide extensive examples in Section 3. All examples are proven in detail with short statements for easy reference.

Our theorems are proven by studying the *intrinsic flat* limit of the manifolds [Definition 29]. This Intinsic flat distance, $d_{\mathcal{F}}(M_1, M_2)$ was originally defined in work of the second author with Wenger [SW11]. It is estimated by explicitly constructing a filling manifold, B^{m+1}, between the two given manifolds, finding the excess boundary manifold A^m satisfying (20) and summing their volumes as in (21). See Remark 2.8 for a straight forward construction. Since $d_{\mathcal{F}}$ depends only on the Riemannian manifolds, M_i, as oriented metric spaces with a notion of integration over m forms, we take settled completions rather metric completions of open domains when analyzing the intrinsic flat distance (see Definition 2.9). If two completely settled oriented Riemannian manifolds, M_1 and M_2 have $d_{\mathcal{F}}(M_1, M_2) = 0$ then there is an orientation preserving isometry between them [SW11]. See Section 2 for a review of the intrinsic flat distance and related concepts.

In Section 4 we prove new explicit estimates on the Gromov-Hausdorff, intrinsic flat and scalable intrinsic flat distances between pairs of manifolds which are diffeomorphic on subdomains [Theorem 4.6]. The subdomains need not be connected. These estimates are found by isometrically embedding the regions into a common metric space defined using a hemispherical construction [Proposition 4.2] and then measuring the Hausdorff, flat and scalable flat distances between their images respectively [Lemma 4.5]. Note that the Hausdorff distance measures distances between the images using tubular neighborhoods while the flat distance measures a filling volume between the images. These estimates have been applied in work of the second author with Dan Lee on questions concerning the Riemannian Penrose Inequality [LSar] and in the first author's doctoral dissertation [Lak].

In Section 5, we prove theorems concerning the intrinsic flat limits of manifolds which converge smoothly away from singular sets. In particular, we prove:

Theorem 1.3. *Let $M_i = (M, g_i)$ be a sequence of oriented Riemannian manifolds such that there is a closed subset, S, of codimension two and a connected precompact exhaustion, W_j, of $M \setminus S$ satisfying (2) such that g_i converge smoothly to g_∞ on each W_j*

$$\operatorname{diam}_{M_i}(W_j) \le D_0 \qquad \forall i \ge j, \tag{6}$$

$$\operatorname{Vol}_{g_i}(\partial W_j) \le A_0, \tag{7}$$

and

$$\operatorname{Vol}_{g_i}(M \setminus W_j) \le V_j \text{ where } \lim_{j\to\infty} V_j = 0. \tag{8}$$

Then

$$\lim_{j\to\infty} d_{\mathcal{F}}(M'_j, N') = 0. \tag{9}$$

where N' is the settled completion of $(M \setminus S, g_\infty)$.

This theorem is a consequence of Theorem 5.2 which assumes only that the connected precompact exhaustion is uniformly well embedded in the sense of Definition 5.1. We discuss the necessities of the conditions for these theorems in Remark 5.3. A key step in the proof is a technical proposition concerning the convergence of exhaustions of manifolds [Proposition 5.4].

In Section 6, we apply the theorems regarding intrinsic flat limits to prove the theorems concerning Gromov-Hausdorff limits mentioned earlier. Note that the second author and Stefan Wenger have proven that the intrinsic flat and Gromov-Hausdorff limits of sequences of manifolds agree when the sequence has nonnegative Ricci curvature and the volume is bounded below uniformly [SW10]. These results are reviewed in Section 6. Theorem 1.3 then immediately implies Theorem 6.6 and Theorem 1.2 when $H = 0$. To obtain Theorem 1.2 for arbitrary values of H, we prove Proposition 6.12.

Applications of these results appear in joint work of the second author and Dan Lee concerning asymptotically flat rotationally symmetric Riemannian manifolds with positive scalar curvature that satisfy an almost equality in the Penrose inequality [LSar]. We believe these results may also be applicable to open questions stated in [LS11]. The first author is examining further applications in his doctoral dissertation.

The authors would like to thank the Simons Center for Geometry and Physics for its hospitality. Attending the many interesting talks there made it clear that a paper clarifying the applications of the intrinsic flat convergence to understand smooth limits away from singular sets would be useful to mathematicians in a wide variety of subfields of geometric analysis.

2. Background

All notions of distances between Riemannian manifolds studied in this paper are built upon Gromov's idea that one may view Riemannian manifolds as metric spaces and isometrically embed them into a common metric space. In this paper, a key part of our work relies on constructing such isometric embeddings. We review Gromov's key ideas in Subsection 2.1.

To estimate the Gromov-Hausdorff distance between a pair of Riemannian manifolds, one needs only find a pair of isometric embeddings $\varphi_i : M_i^m \to Z$ into a common complete metric space Z and then measure the Hausdorff distances between the images. We review the definition of the Hausdorff and Gromov-Hausdorff in Subsection 2.2.

To estimate the Sormani-Wenger intrinsic flat distance one must measure the flat distance between these images. So one may construct a Riemannian manifold of one dimension higher filling in the space between the two images, possible with some excess boundary. Note that one can only measure the intrinsic flat distance between oriented manifolds with finite volume of the same dimension. See Subsection 2.3.

The scalable intrinsic flat distance is also defined using filling manifolds and excess boundaries. It is reviewed in Subsection 2.4.

Remarks 2.3, 2.6, 2.8 and 2.11 capture the key properties of these three notions of distance needed to estimate them for the purposes of this paper.

2.1. **Metric Spaces and Isometric Embeddings.**

Definition 2.1. *Recall that one may view a Riemannian manifold* (M, g) *as a metric space* (M, d) *by defining the distances between points as follows:*

$$d(x_1, x_2) = \inf \left\{ L_g(\gamma) : \gamma(0) = x_1, \gamma(1) = x_2 \right\} \tag{10}$$

where

$$L_g(\gamma) = \int_0^1 g(\gamma'(t), \gamma'(t))^{1/2} dt \tag{11}$$

Given a connected subdomain, $W \subset M$*, and* $x, y \in W$*, the "restricted metric",* $d_M(x, y)$*, will denote the distance between* x *and* y *measured as in (10) where the infimum taken over all curves* $\gamma : [0, 1] \to M$*, while the "induced length metric",* $d_W(x, y) \geq d_M(x, y)$*, has the infimum taken only over curves* $\gamma : [0, 1] \to W$*. We denote the restricted and intrinsic length diameters of* $U \subset W \subset M$ *as follows*

$$\operatorname{diam}_M(U) = \sup\{d_M(x, y) : x, y \in U\} \tag{12}$$

$$\operatorname{diam}_W(U) = \sup\{d_W(x, y) : x, y \in U\} \tag{13}$$

More generally a length metric space is a metric space whose distances are defined as an infimum of lengths of rectifiable curves. Compact length metric spaces always have minimizing geodesics between points achieving the distance.

In this paper we will often define metric spaces, Z, by gluing together Riemannian manifolds with corners along their boundaries. In this way we may still apply (10) to define the distances between points. Again, for connected subdomains, $W \subset Z$, one has both an induced length metric, d_W, and a restricted distance $d_Z \leq d_W$ just as in Definition 2.1.

Definition 2.2. *An isometric embedding $\varphi : X \to Z$ is a distance preserving map*

$$d_Z(\varphi(x_1), \varphi(x_2)) = d_X(x_1, x_2) \qquad \forall x_1, x_2 \in X \tag{14}$$

One should be aware that a Riemannian isometric embedding defined by the fact that $d\varphi$ is an isometry on the tangent spaces at each point, is not necessarily an isometric embedding. For example, the natural embedding of the sphere into Euclidean space is not an isometric embedding with the standard metric on the sphere. See Figure 1.



FIGURE 1. S^1 in the center isometrically embeds into S^2 on the right, but does not isometrically embed into $\mathbb{E}^2$ on the left.

Remark 2.3. *Suppose two manifolds, M_i have diffeomorphic subdomains, U_i, then a filling manifold can be constructed of the form $U \times [h_1, h_2]$ with a well chosen metric g' so that M_i isometrically embed into*

$$Z = M_1 \sqcup (U \times [h_1, h_2]) \sqcup M_2. \tag{15}$$

Here Z is glued together so that U_i is identified point to point with $U \times \{h_i\}$. A precise way of choosing such a g' will be given in Theorem 4.6. See Figure 2.

2.2. **The Gromov-Hausdorff Distance.** The Gromov-Hausdorff distance between a pair of Riemannian manifolds is estimated by taking isometric embeddings into a common metric space Z and measuring the Hausdorff distance between them. This distance was introduced by Gromov in [Gro99]. It is defined on pairs of metric spaces.

Definition 2.4 (Hausdorff). *Given two subsets $Y_1, Y_2 \subset Z$, the Hausdorff distance is defined*

$$d_H^Z(Y_1, Y_2) = \inf\{r : Y_1 \subset T_r(Y_2) \text{ and } Y_2 \subset T_r(Y_1)\} \tag{16}$$

where $T_r(Y) = \{z \in Z : \exists y \in Y \text{ s.t. } d(y,z) < r\}$.

One may immediately observe that the topology and dimension of subsets which are close in the Hausdorff sense can be quite different.

Definition 2.5 (Gromov). *Given a pair of metric spaces (X_1, d_1) and (X_2, d_2), the Gromov-Hausdorff distance between them is*

$$d_{GH}(X_1, X_2) = \inf\left\{d_H^Z(\varphi_1(X_1), \varphi_2(X_2)) : \varphi_i : X_i \to Z\right\} \tag{17}$$

where the infimum is taken over all common metric spaces, Z, and all isometric embeddings, $\varphi_i : X_i \to Z$.



Figure 2. M_1 and M_2 depicted on the left and the right isometrically embed into Z in the center. See Remarks 2.3 and 2.6.

Remark 2.6. *In Figure 2 depicting Remark 2.3, we see that*

$$d_{GH}(M_1, M_2) \le d_H^Z(\varphi_1(M_1), \varphi_2(M_2)) \tag{18}$$

which is roughly the length of a curve from the tip of the bump in M_1 running back within $\varphi(M_1) \subset Z$ to the warped region $U \times [h_1, h_2]$ and then straight up to M_2. Later in this paper Theorem 4.6 we will find a precise description of the metric on the metric space Z of Figure 2.

Gromov proved in [Gro99] that this a distance between compact metric spaces, in the sense that $d_{GH}(X_1, X_2) = 0$ iff X_1 and X_2 are isometric. In general one takes the metric completion, $\bar{X}$, of a precompact space, X, before discussing it's Gromov-Hausdorff distance and we will do the same here. Recall that the metric completion is defined as follows:

Definition 2.7. *Given a precompact metric space, X, the metric completion, $\bar{X}$ of X is the space of Cauchy sequences, $\{x_j\}$, in X with the metric*

$$d(\{x_j\},\{y_j\}) = \lim_{j\to\infty} d_X(x_j,y_j) \tag{19}$$

and where two Cauchy sequences are identified if the distance between them is 0. *There is an isometric embedding, $\varphi : X \to \bar{X}$, defined by $\varphi(x) = \{x\}$ where $\{x\}$ is a constant sequence. Lipschitz functions, $F : X \to Y$, extend to $F : \bar{X} \to Y$ via $F(\{x_j\}) = \lim_{j\to\infty} F(x_j)$ as long as Y is complete.*

Gromov's compactness theorem states that a sequence of Riemannian manifolds M^m_j with a uniform lower bound on Ricci curvature have a subsequence which converges in the Gromov-Hausdorff sense. More generally one may replace the Ricci curvature bound with a bound on the number, $N(r)$, of disjoint balls of radius, r, that can be placed in a metric space, X. That is, a sequence of metric spaces X_j with a uniform bound on $N(r)$ for all r sufficiently small, has a subsequence which converges in the Gromov-Hausdorff sense to a compact limit space X. Conversely, if $d_{GH}(X_j, X) \to 0$, then there is a uniform bound on $N(r)$. [Gro99].

2.3. **The Intrinsic Flat Distance.**

To estimate the intrinsic flat distance between a pair of oriented Riemannian manifolds one again needs only find a pair of isometric embeddings, $\varphi_i : M^m_i \to Z$, into a common complete metric space, Z. When one finds a filling submanifold, $B^{m+1} \subset Z$, and an excess boundary submanifold, $A^m \subset Z$, such that

$$\int_{\varphi_1(M_1)} \omega - \int_{\varphi_2(M_2)} \omega = \int_B d\omega + \int_A \omega, \tag{20}$$

then the intrinsic flat distance is bounded by

$$d_{\mathcal{F}}(M^m_1, M^m_2) \le \mathrm{Vol}_m(A^m) + \mathrm{Vol}_{m+1}(B^{m+1}). \tag{21}$$

Generally the filling manifold and excess boundary can have corners and more than one connected component. See Figure 7.

Remark 2.8. *In Figure 2 depicting Remark 2.3, we have M_i isometrically embedded into a well chosen metric space*

$$Z = M_1 \sqcup (U \times [h_1,h_2]) \sqcup M_2. \tag{22}$$

Applying (20) to

$$B = U \times [h_1,h_2] \tag{23}$$

we see that the excess boundary

$$A = (M_1 \setminus U_1) \cup (\partial U \times [h_1,h_2]) \cup (M_2 \setminus U_2). \tag{24}$$

Then

$$\begin{aligned} d_{\mathcal{F}}(M_1, M_2) &\leq \operatorname{Vol}_m(M_1 \setminus U_1) + \operatorname{Vol}_m(M_2 \setminus U_2) + \operatorname{Vol}_m(\partial U \times [h_1, h_2]) \\ &+ \operatorname{Vol}_{m+1}(U \times [h_1, h_2], g'). \end{aligned}$$

An explicit construction of the metric g' on $U \times [h_1, h_2]$ in Theorem 4.6, allows one to precisely estimate the volume of $U \times [h_1, h_2]$ and $\partial U \times [h_1, h_2]$.

To understand limits of sequences of Riemannian manifolds, the intrinsic flat distance was defined on a larger class of metric spaces called integral current spaces in [SW11]. An integral current space, (X, d, T), is a metric space, X, with a metric, d, and an integral current structure, T, such that set(T) = X. An oriented Riemannian manifold, (M, g), of finite volume, has a metric, d_M, defined as in Definition 2.1 and an integral current structure, T, acting on m dimensional differential forms, ω as

$$T(\omega) = \int_M \omega. \tag{25}$$

More generally, the integral current structure, T, of an integral current space, (X, d, T), is an m dimensional integral current $T \in \mathbf{I}_m(\bar{X})$ defined as in Ambrosio-Kirchheim's work [AK00]. The integral current structure T provides both an orientation and a measure called the mass measure denoted $\|T\|$ and set(T) is the set of positive lower density for this measure. On a oriented Riemannian manifold, the mass measure is just the Lebesgue measure. More generally the mass measure can have integer valued weights.

If (M^m, g) is a Riemannian manifold with singularities on a subset S and the Hausdorff measure, $\mathcal{H}_m(S) = 0$, then one obtains a corresponding integral current space by taking the settled completion of $M \setminus S$ defined as follows:

Definition 2.9. [SW11] *The settled completion, X', of a metric space X with a measure μ is the collection of points x in the metric completion $\bar{X}$ which have positive lower density:*

$$\liminf_{r \to 0} \mu(B_p(r))/r^m > 0. \tag{26}$$

The resulting space is then "completely settled".

If a manifold has only point singularities, one includes all conical tips and does not include cusp tips in a manifold with point singularities. See Figure 3. This is natural because the essential property of an integral current space is its integration and points of 0 lower density do not contribute to that integration. In fact integral current spaces are completely settled with respect to the mass measure, $\|T\|$, as a consequence of the requirement that set(T) = X.

The mass of an integral current space, $\mathbf{M}(T)$, is a weighted volume of sorts which takes into account the integer valued Borel weight defining the current structure on



FIGURE 3. The completion, $\bar{X}$ includes the boundary and fills in the three "holes" and the settled completion, X', removes the cusped singularity but keeps the boundary, cone tip and smoothly filled hole.

the space. When the integral current space is an oriented Riemannian manifold then its mass is just its volume, $\mathbf{M}(M) = \mathrm{Vol}(M)$.

The boundary of an integral current space is defined

$$\partial(X, d, T) = (\mathrm{set}(\partial \mathrm{T}), \mathrm{d}, \partial \mathrm{T}) \tag{27}$$

where ∂T is the boundary of the integral current defined as in [AK00] so that it satisfies Stoke's Theorem. When M is a Riemannian manifold, its boundary is just the usual boundary, ∂M.

The flat distance between two integral currents is defined as in [FF60]

$$d_F(T_1, T_2) = \inf\left\{\mathbf{M}(B^{m+1}) + \mathbf{M}(A^m) : \, T_1 - T_2 = A + \partial B\right\}. \tag{28}$$

That is $T_1(\omega) - T_2(\omega) = A(\omega) + \partial B(\omega) = A(\omega) + B(d\omega)$.

Definition 2.10 (Sormani-Wenger)**.** *The intrinsic flat distance between integral current spaces is defined in* [SW11] *as*

$$d_{\mathcal{F}}((X_1, d_1, T_1), (X_2, d_2, T_2)) = \inf\left\{d_F^Z(\varphi_{1\#}T_1, \varphi_{2\#}T_2);\ \varphi_i : X_i \to Z\right\} \tag{29}$$

where the infimum is taken over all common complete metric spaces, Z, and all isometric embeddings $\varphi_i : X_i \to Z$ and where $\varphi_\#$ is the push forward map on integral currents.

If one constructs a specific Z and isometric embeddings $\varphi_i : M_i \to Z$, then one needs only estimate the flat distance between the images to obtain an upper bound for the infimum in (29). An explicit filling manifold B satisfying (20), then provides an upper bound on the infimum in (28). This is how one obtains the estimate in (21). See also Remark 2.8.

In [SW11] it is proven that this is a distance between precompact integral current spaces in the sense that $d_{\mathcal{F}}((X_1, d_1, T_1), (X_2, d_2, T_2)) = 0$ iff there is a current preserving isometry from X_1 to X_2. When the integral current spaces are oriented manifolds, then there is an orientation preserving isometry.

Note that all integral current spaces are metric spaces but they need not be length spaces. As will be seen in Example 3.4 a sequence of connected Riemannian manifolds may converge in the intrinsic flat sense to an integral current space which has broken apart due to the development of a cusp singularity. So the limit is not a length metric space.

In [SW11] it is proven that if (M, g_j) converge smoothly to (M, g_∞) then they converge in the intrinsic flat sense. In fact, precise estimates on the intrinsic flat distance are given in terms of the Lipschitz distance, the diameters and the volumes of the spaces. The bounds are found using geometric measure theory. Here we provide a new estimate relating the intrinsic flat and Lipschitz distances by explicitly constructing a filling manifold between them [Lemma 4.5].

In [SW11] it was proven that if a sequence of oriented Riemannian manifolds with a uniform upper bound on their volumes and volumes of their boundaries converges in the Gromov-Hausdorff sense to a compact metric space (Y, d), then a subsequence converges in the intrinsic flat sense to (X, d, T) where $X \subset Y$ and the metric d is restricted from Y. In Example 3.4 we see that this may be a proper subset. In fact the Intrinsic flat limit may be the $(0, 0, 0)$ integral current space if (Y, d) has a lower dimension than the manifolds in the sequence [SW11].

In [SW10] two theorems were proven indicating when the intrinsic flat limit and the Gromov-Hausdorff limits agree. These theorems will be reviewed later in the paper as they are applied. We will also apply the techniques in their proofs to prove Theorem 1.2.

2.4. **The Scalable Intrinsic Flat Distance.** The scalable intrinsic flat distance was suggested as a notion in work of the second author with Dan Lee [LS11] following a recommendation of Lars Andersson. It is defined to scale with distance when the Riemannian manifolds are rescaled. In particular,

$$d_{s\mathcal{F}}(M_1^m, M_2^m) \le (\mathrm{Vol}_m(A^m))^{1/m} + \left(\mathrm{Vol}_{m+1}(B^{m+1})\right)^{1/(m+1)} \tag{30}$$

whenever there exist isometric embeddings, $\varphi_i : M_i^m \to Z$, into a common complete metric space, Z, and one finds a filling submanifold, $B^{m+1} \subset Z$, and an excess boundary submanifold, $A^m \subset Z$, satisfying (20).

Remark 2.11. *In the setting of Remark 2.8 depicted in Figure 2 we see that*

$$\begin{aligned} d_{s\mathcal{F}}(M_1, M_2) \;\; &\le \;\; \mathrm{Vol}_{m+1}(U \times [0, h], g')^{1/(m+1)} \\ &+ \;\; (\mathrm{Vol}_m(M_1 \setminus U_1) + \mathrm{Vol}_m(M_2 \setminus U_2) + \mathrm{Vol}_m(\partial U \times [h_1, h_2], g'))^{1/m} . \end{aligned}$$

More precisely:

Definition 2.12. *The scalable intrinsic flat distance between integral current spaces is defined as*

$$d_{s\mathcal{F}}((X_1, d_1, T_1), (X_2, d_2, T_2)) = \inf\left\{ d^Z_{sF}(\varphi_{1\#}T_1, \varphi_{2\#}T_2);\ \varphi_i : X_i \to Z \right\} \tag{31}$$

where the infimum is taken over all common complete metric spaces, Z, and all isometric embeddings $\varphi_i : X_i \to Z$ and where $\varphi_\#$ is the push forward map on integral currents and where the scalable flat distance between m dimensional integral currents is defined by

$$d_{sF}(T_1, T_2) = \inf\left\{\mathbf{M}(B)^{1/(m+1)} + \mathbf{M}(A)^{1/m} : T_1 - T_2 = A + \partial B\right\} \tag{32}$$

This distance is being studied in more detail in work of Jorge Basilio [Bas].

3. Examples

The following examples are presented to indicate how little control one may have on limits of manifolds which converge smoothly away from singular sets and to prove the necessity of the conditions in our theorems. The proofs of these examples will sometimes rely on our theorems proven below but we include them up front so that they can be kept in mind when reading the remainder of the paper.

3.1. **Losing a Region.**

Example 3.1. *There are metrics, g_j, on the sphere, M^3, such that (M^3, g_j) converge smoothly away from $S = \bar{B}_{p_0}(\pi/16)$, such that the metric completion of the smooth limit away from S is $S^3 \setminus B_{p_0}(\pi/16)$, the standard round sphere (S^3, g_0) with a ball removed. The smooth limit of M^3 without the singular set removed is the entire round sphere and this agrees with the intrinsic flat and Gromov-Hausdorff limits (c.f. Lemma 4.5).*

Proof. Taking the metrics, $g_j = g_0$, we have a constant sequence of standard spheres. So the intrinsic flat and Gromov-Hausdorff limits are clearly the standard sphere. Furthermore $(g_j, M \setminus S)$ clearly converges to $(g_0, M \setminus \bar{B}_{p_0}(\pi/16))$ whose metric completion is $(g_0, M \setminus B_{p_0}(\pi/16))$. □

3.2. **Cones and Cusps.**

Example 3.2. *There are metrics g_j on the sphere M^3 such that (M^3, g_j) converge smoothly away from a point singularity $S = \{p_0\}$ and the metrics g_j form a conical singularity at p_0. The Gromov-Hausdorff and intrinsic flat limits agree with the metric completion of $(M \setminus S, g_\infty)$ which is the sphere including the conical tip.*

Proof. More precisely the metrics g_j are defined by

$$g_j = dr^2 + f_j^2(r) g_{S^2} \text{ for } r \in [0, \pi] \tag{33}$$

where $f_j(r) = (1/j)\sin(r) + (1 - 1/j) f(r)$ in which, $f(r)$ is a smooth function such that:

$$f(r) = \sin(r) \text{ for } r \in [0, \pi/2], \tag{34}$$

and,

$$f(r) = -\frac{2}{\pi}(r-\pi) \text{ for } r \in [3\pi/4, \pi]. \tag{35}$$

For any $\delta > 0$, f_j converge to f smoothly on $[0, \pi - \delta]$. Thus g_j converge smoothly on compact subsets of $M \setminus S$ to

$$g_\infty = dr^2 + f^2(r) g_{S^2}. \tag{36}$$

The metric completion of $(M \setminus S, g_\infty)$ then adds in a single point p_0 at $r = \pi$. Since

$$\liminf_{r\to 0} \mu(B_{p_0}(r))/r^3 = \frac{4}{3\pi^2} vol(S^2) = \frac{16}{3\pi} > 0, \tag{37}$$

the point, p_0, is also included in the settled completion of $(M \setminus S, g_\infty)$. To complete the proof of the claim we could apply Theorem 1.3. □

Example 3.3. *There are metrics g_j on the sphere M^3 such that (M^3, g_j) converge smoothly away from a point singularity $S = \{p_0\}$ and the metrics g_j form a cusp singularity at p_0. The Gromov-Hausdorff agree with the metric completion of $(M \setminus S, g_\infty)$ which is the sphere including the cusped tip. However the intrinsic flat limit of $(M \setminus S, g_\infty)$ does not include the cusped tip because it has 0 density. So the intrinsic flat limit is the settled completion of $(M \setminus S, g_\infty)$ which in this case is $(M \setminus S, g_\infty)$*

Proof. More precisely the metrics g_j are defined by

$$g_j = dr^2 + f_j^2(r) g_{S^2} \text{ for } r \in [0, \pi] \tag{38}$$

where $f_j(r) = (1/j)\sin(r) + (1 - 1/j) f(r)$ in which, $f(r)$ is a smooth function such that:

$$f(r) = \sin(r) \text{ for } r \in [0, \pi/2], \tag{39}$$

and,

$$f(r) = \frac{4}{\pi^2}(r-\pi)^2 \text{ for } r \in [3\pi/4, \pi]. \tag{40}$$

For any $\delta > 0$, f_j converge to f smoothly on $[0, \pi - \delta]$. Thus g_j converge smoothly on compact subsets of $M \setminus S$ to

$$g_\infty = dr^2 + f^2(r) g_{S^2}. \tag{41}$$

The metric completion of $(M \setminus S, g_\infty)$ then adds in a single point p_0 at $r = \pi$. Since

$$\liminf_{r\to 0} \mu(B_{p_0}(r))/r^3 = \liminf_{r\to 0} \frac{4}{5\pi^2} r^2 vol(S^2) = 0, \tag{42}$$

the point, p_0, is not included in the settled completion of $(M \setminus S, g_\infty)$.

This Gromov-Hausdorff and Intrinsic Flat limits in this example were proven to be as claimed in the Appendix of [SW11]. One may also apply Theorem 4.6 to reprove this. □

3.3. Not Connected.

Example 3.4. *There are smooth metrics g_i on the sphere, M^3, converging smoothly away from the equator, S, such that the equator pinches to* 0*. Then $(M^3 \setminus S, g_i)$ has two components, each converging to a standard sphere with a point removed. The metric completion of each of the two disjoint metric spaces is a standard sphere. However the Gromov-Hausdorff limit is a pair of spheres joined at a point singularity. So we see why connectedness of the singular set is a necessary condition in Theorem 5.2. Here the singular set is codimension 1.*

Remark 3.5. *In upcoming work of the first author* [Lak]*, appropriate gluings of disjoint metric spaces are taken to recover the Gromov-Hausdorff limit when $M \setminus S$ is not connected.*

Proof. We take the sequence of metrics

$$g_i = dr^2 + f_i^2(r) g_{S^2} \tag{43}$$

where $f_i = \frac{1}{i}\sin(r) + \frac{i-1}{2i}|\sin(2r)|$. These are smooth metrics for $r \in [0,\pi]$ because $f_i(r) > 0$ for $r \in (0,\pi)$,

$$f_i'(0) = \frac{1}{i} + \frac{2(i-1)}{2i} = 1, \tag{44}$$

$$f_i'(\pi) = -\frac{1}{i} - \frac{2(i-1)}{2i} = -1 \tag{45}$$

and $f_i''(0) = f_i''(\pi) = 0$. As $i \to \infty$, g_i converge smoothly away from $r^{-1}(\pi/2)$ to

$$g_\infty = dr^2 + \frac{\sin^2(2r)}{4} g_{S^2} \tag{46}$$

which is a metric on a pair of spheres, each with a point removed. The metric completion keeps the pair of spheres disjoint, endowing each with its own point of completion.

The Gromov-Hausdorff and intrinsic flat limits however are a connected pair of spheres joined at point which creates a conical singularity. This can be seen because in fact the distances d_i defined on M_i using g_i are in fact converging in the Lipschitz sense to d_∞ defined using the infimum of lengths, L_∞, of curves between points where

$$L_\infty(C) = \int_0^1 g_\infty(C'(s), C'(s))^{1/2}\, ds. \tag{47}$$

Taking $W_j = r^{-1}[0, \pi/2 - 1/j] \cup r^{-1}[\pi/2 + 1/j, \pi]$, then we have smooth convergence on W_j. The uniform embeddedness constants converge to 0. Both $\mathrm{Vol}_{g_i}(V \setminus Wj) < V_j$ with $V_j \to 0$ and $\mathrm{Vol}_{q_i}(\partial W_j) \leq A_j$ with $A_j \to 0$. So we only fail the connectedness hypothesis of this theorem. □

3.4. **Bubbling.**

Example 3.6. *There are smooth metric g_i on the sphere, M^3 converging smoothly away from the singular set $S = \{p_0\}$ to a sphere. Yet (M^3, g_i) converge in the Gromov-Hausdorff and intrinsic flat sense to a pair of spheres meeting at p_0. See Figure 4.*



FIGURE 4. Example 3.6.

Proof. Let

$$g_i = h_i^2(r)dr^2 + f_i^2(r)g_{S^2} \tag{48}$$

where

$$h_i(r) = 2 \quad \text{on} \quad r \in [0, a_i] \tag{49}$$

$$f_i(r) = \sin(r) \quad \text{on} \quad r \in [0, a_i] \tag{50}$$

(51)

where $a_i = \pi - \pi/(10i)$ so that g_i converges smoothly away from S to the round metric g_∞ on the sphere. The metric completion of $(M \setminus S, g_\infty)$ is the round sphere.

Now we set

$$h_i(r) = 10i \quad \text{on} \quad r \in [b_i, \pi] \tag{52}$$

$$f_i(r) = \sin((r-\pi)/(10i)) \quad \text{on} \quad r \in [b_i, \pi] \tag{53}$$

(54)

where $b_i = \pi - (\pi - \pi/(10i))/(10i)$ so there is symmetry and extend them smoothly for $r \in [a_i, b_i]$ so that

$$h_i(a_i) = 2 \le h_i(r) \le 10i = h_i(b_i) \tag{55}$$

and

$$0 < f_i(r) < \max\{f_i(a_i), f_i(b_i)\}. \tag{56}$$

These thin regions are converging to a single point. So the Gromov-Hausdorff limit of (M, g_i) is a pair of standard spheres joined at a point and the intrinsic flat limit is the same. The smooth limit away from S missed the second sphere! □

3.5. **Losing Volume in the Limit.**

Example 3.7. *There are (M^3, g_i) all isometric to the standard sphere which converge smoothly away from a singular set $S = \{p_0\}$ to $(M \setminus S, g_\infty)$ which is isometric to an open hemisphere. The metric completion agrees with the settled completion, (M_∞, d_∞) which is isometric to a closed hemisphere. The singular set is codimension 2 in M. This example satisfied all the conditions of all of our Theorems concerning smooth convergence away from singular sets except* $\mathrm{Vol}(M \setminus W_j) < V_j$ *where* $\lim_{j\to\infty} V_j = 0$.

Proof. Again we view $M^3 = S^3$ as a warped product with a warping function $r \in [0, \pi]$, such that $r(p_0) = \pi$. Let

$$g_i = (h_i'(r))^2 dr^2 + \sin^2(h_i(r)) g_{S^2} \tag{57}$$

where $h_i(r)$ is a smooth increasing function such that

$$h_i(r) = r(\pi/2)/(\pi - \pi/(2i)) \text{ for } r \in [0, \pi - 1/i] \tag{58}$$

$$h_i(r) = \pi - (r - \pi)(1/(2i))(\pi - 1/(2i)) \text{ for } r \in [\pi - 1/(2i), \pi]. \tag{59}$$

Then the diffeomorphism which maps $r \mapsto s = h_i(r)$ is an isometry from (M^3, g_i) to (S^3, g_{S^3}).

On any compact set $K \subset M \setminus S$, there exists a j sufficiently large that $K \subset r^{-1}[0, \pi - 1/j]$. Taking $i \to \infty$ we see that on K, $h_i(r) \to r/2$ and g_i converge to

$$g_\infty = (1/2)^2 dr^2 + \sin^2(r/2) g_{S^2}. \tag{60}$$

Thus $(M \setminus S, g_\infty)$ is isometric to an open hemisphere via the isometry which maps $r \mapsto s = r/2$. The metric completion is then the closed hemisphere and the settled completion agrees with the metric completion because every point in the closed hemisphere has positive lower density.

Setting $W_j = r^{-1}[0, \pi - 1/j]$, we see that

$$\mathrm{Vol}_{g_i}(\partial W_j) \le 4\pi. \tag{61}$$

Clearly the diameter, volume, Ricci curvature and contractibility conditions all hold because the sequence of (M, g_j) are all isometric to spheres. However

$$\lim_{i\to\infty} \mathrm{Vol}_{g_i}(M \setminus W_j) \ge \mathrm{Vol}(S^3)/2. \tag{62}$$

□

3.6. **Unbounded Volumes.**

Example 3.8. *There are metrics g_i on the sphere M^3 with nonnegative Ricci curvature such that (M^3, g_i) converge smoothly away from a point singularity $S = \{p_0\}$ to a complete noncompact manifold; In particular, converging to a hemisphere attached to a cylinder of length k on the $r^{-1}[0, \pi - 1/k)$ region. This example satisfies all the hypothesis of Theorem 1.2 except (4).*

Proof. For any $L \in \mathbb{R}$ large enough, define the warped metric g_L on $[0, L] \times S^2$ as follows:

$$g_L(t) = dt^2 + (f_L(t))^2 g_{S^2} \tag{63}$$

where,

$$f_L(t) = \sin(t) \text{ for } t \in [0, \pi/2] \tag{64}$$

$$f_L(t) = 1 \text{ for } t \in [\pi/2 + 1/100, t - \pi/2 - 1/100] \tag{65}$$

$$f_L(t) = \sin(\pi + t - L) \text{ for } t \in [L - \pi/2, L] \tag{66}$$

and $f_L(t)$ smooth with $f_L"(t) < 0$ elsewhere. We will be calling g_L, the double torpedo metric (it is comprised of two torpedo metrics glued together from their cylindrical ends.) For any L, g_L has nonnegative Ricci curvature.

Let $\phi : [0, \pi] \to [0, \infty)$ be a smooth increasing function such that

$$\phi(r) = r \text{ for } r \in [0, \pi/2] \tag{67}$$

with

$$\lim_{r \to \pi} \phi(r) = \infty \tag{68}$$

For $j > 2$, let $\phi_j(r) : [0, \pi] \to [0, L_j = j + \pi/2 + 1]$ be a smooth increasing function such that

$$\phi_j(r) = \phi(r) \text{ for } r \in [0, \phi^{-1}(j + \pi/2)], \tag{69}$$

and

$$\phi_j(r) = j + r - \pi/2 + 1 \text{ for } r \text{ near } \pi. \tag{70}$$

Again, we view $M^3 = S^3$ as a warped product with a warping function $r \in [0, \pi]$, such that $r(p_0) = \pi$. Let

$$g_j(r) = \phi_j^*\left(g_{L_j}\right) = (\phi_j'(r))^2 dr^2 + \left(f_{L_j}\right)^2 (\phi_j(r)) g_{S^2} \tag{71}$$

Then the diffeomorphism ϕ_j is an isometry from (M^3, g_j) to (S^3, g_{L_j}). On any compact set $K \subset M \setminus S$, there exists a k sufficiently large that $K \subset r^{-1}[0, \pi - 1/k]$. Taking $j \to \infty$ we see that on K, g_j converge to

$$g_\infty = (\phi'(r))^2 dr^2 + f^2(\phi(r)) g_{S^2}. \tag{72}$$

where,

$$f(r) = \sin(r) \;\text{ for }\; r \in [0, \pi/2 - 1/100] \;\text{ and }\; f(r) = 1 \;\text{ for }\; r \in [\pi/2 + 1/100, \infty) \tag{73}$$

which is a hemisphere smoothly attached to a cylinder of length k.

If we take $W_j = r^{-1}([0, \pi - 1/j))$ then, we see that, $\mathrm{Vol}_{g_i}(W_j)$ and $\mathrm{Vol}_{g_i}(M \setminus W_j)$ are unbounded. $(M \setminus S, g_\infty)$ is complete hence coincides with the metric completion. Since $(M \setminus S, g_\infty)$ is noncompact , (M^3, g_j) does not have Gromov - Hausdorff limit. Also since the volume is not finite, there is no intrinsic flat limit either.

Nevertheless, this example has $\mathrm{Vol}(\partial W_j) \leq 4\pi$ and W_j are uniformly embedded, the sequence has nonnegative Ricci curvature and a uniform contractibility function, $\rho(r) = r$ for $r \in (0, \pi/2]$. □

3.7. **Spheres with Splines.** The following examples are based upon examples in [SW11].

Example 3.9. *There are metrics g_i on the sphere M^3 such that (M^3, g_i) converge smoothly away from a point singularity $S = \{p_0\}$ yet we have a single spline of finite length, L, becoming thinner and thinner so that the Gromov-Hausdorff limit is not the sphere while the intrinsic flat limit is just the sphere. The metric completion of $(M \setminus S, g_\infty)$ is also the sphere in this example.*

A version of this example with positive scalar curvature will be given in [Lak].

Proof. More precisely the metrics g_i are defined by

$$g_i = h_i(r)^2 dr^2 + f_i^2(r) g_{S^2} \text{ for } r \in [0, \pi] \tag{74}$$

where $f_i(r) = \sin(r)$ and

$$h_i(r) = 1 + i \exp\left(\frac{\left(\frac{1}{2i}\right)^2}{\left(r - \pi + \frac{2}{i}\right)\left(r - \pi + \frac{1}{i}\right)} \right) \chi_{\left[\pi - \frac{2}{i}, \pi - \frac{1}{i})\right]}. \tag{75}$$

Observe that on $r^{-1}[0, \pi - 1/j)$ we have $h_i(r) = 1$ for $i \geq 2j$. So g_i converge smoothly away from p_0 to the standard metric on a sphere, g_∞. The metric and settled completions of $(M \setminus \{p_0\}, g_\infty)$ are both the standard sphere.

We will refer to $N_i = r^{-1}(\pi - 2/i, 0]$ with the metric g_i as a spline. Observe that

$$\mathrm{diam}_{g_i}(M_j) \geq \int_0^\pi h_i(r)\, dr = \pi - 2/i + L + 1/i \tag{76}$$

where L is the length of the spline:

$$L = \int_{\pi - 2/i}^{\pi - 1/i} h_i(r)\, dr \tag{77}$$

$$= \int_0^1 e^{1/(4u(u-1))}\, du \tag{78}$$

Since the diameter of the Gromov-Hausdorff limit, when it exists, is the limit of the diameters of the sequence, we see that the Gromov-Hausdorff limit is not metric completion in this case. We will not provide an explicit proof that the Gromov-Hausdorff limit is in fact the sphere iwth a line segment of length D attached at p_0.

Now taking $W_j = r^{-1}([0, \pi - 1/(2j)])$, we see that

$$\mathrm{diam}_{M_i}(W_j) \leq \pi \text{ for } i \geq j, \tag{79}$$

$$\mathrm{Vol}(M_i) \leq \mathrm{Vol}(S^3, g_0) + \sin(1/(2i))L \leq V_0 \tag{80}$$

$$\mathrm{Vol}_{g_i}(\partial W_j) \leq 4\pi 1 \tag{81}$$

$$\mathrm{Vol}_{g_i}(M \setminus W_j) \leq \pi(1/(2j)^2) + \pi\sin(2/j)^2 L \leq V_j \tag{82}$$

where $\lim_{j\to\infty} V_j = 0$. By Theorem 1.3 we have the intrinsic flat limit is settled metric completion which is the sphere. This example has no uniform lower bound on Ricci curvature nor a uniform contractibility function so it demonstrates the necessities of these conditions in all of our theorems which require them to prove the Gromov-Hausdorff limit exists and is the metric completion of $(M \setminus S, g_\infty)$. □

Example 3.10. *There are metrics g_i on the sphere M^3 with uniformly bounded diameter and volume such that (M^3, g_i) converge smoothly away from a point singularity $S = \{p_0\}$ and we have increasingly many splines of length L whose total volume goes to 0 based in smaller and smaller neighborhoods of S. The metric completion of $(M \setminus S, g_\infty)$ is the round sphere. This is also the intrinsic flat limit. The Gromov-Hausdorff limit, however, does not exist since the number of balls of radius $L/2$ diverges to infinity.*

A version of this example with positive scalar curvature will be given in [Lak].

Proof. Let (M, g_i) be created by taking the standard sphere of radius 1 and removing i pairwise disjoint balls of radius $2/i^2$ from the ball of radius $2/i$ about p_0. Replace each of those balls with a spline, N_{i^2} from the previous example. Each spline has length L as in the previous example, so there are i balls of radius $L/2$ centered at the tips of the splines. By Gromov's Compactness Theorem's Converse, there is no subsequence converging in the Gromov-Hausdorff sense.

However, $\mathrm{diam}(M, g_i) \leq \pi + 2L$.

Each (M, g_i) is diffeomorphic to S^3, via the identity map outside of the splines and via the diffeomorphism from each spline to the ball it has replaced. Taking any precompact set $W \subset S^3$, such that $p_0 \notin \bar{W}$, we can take i sufficiently large that $W \cap B_{p_0}(2i) = \emptyset$, so that g_i is then the standard metric on W_i. So we see that (M, g_i) converges smoothly to a standard sphere with p_0 removed. The metric and settled completions are one again the standard sphere.

Let $W_j = S^3 \setminus B_{p_0}(2j)$ where the ball is measured using the standard metric on S^3 so that for $j \geq i$ there are no splines within (W_j, g_i).

$$\operatorname{diam}_{M_i}(W_j) \leq \pi \text{ for } i \geq j, \tag{83}$$

$$\operatorname{Vol}(M_i) \leq \operatorname{Vol}(S^3, g_0) + i\sin(1/(2i^2))L \leq V_0 \tag{84}$$

$$\operatorname{Vol}_{g_i}(\partial W_j) \leq 4\pi\sin(1/2i^2)i \leq A_0 \tag{85}$$

$$\operatorname{Vol}_{g_i}(M \setminus W_j) \leq \pi i(1/(2i^2)^2) + \pi i\sin(2/i^2)^2 L \leq V_j \tag{86}$$

where $\lim_{j\to\infty} V_j = 0$. By Theorem 1.3 we have the intrinsic flat limit is settled metric completion which is the sphere. This example has no uniform lower bound on Ricci curvature nor a uniform contractibility function so it demonstrates the necessities of these conditions in all of our theorems which require them to prove the Gromov-Hausdorff limit exists and is the metric completion of $(M \setminus S, g_\infty)$. □

Example 3.11. *There are metrics g_i on the sphere M^3 s with uniformly bounded volume such that (M^3, g_i) converge smoothly away from a point singularity $S = \{p_0\}$ and we have a single spline of increasing length whose volume goes to 0 and width goes to 0 contained in smaller and smaller neighborhoods of S. The metric completion of $(M \setminus S, g_\infty)$ is the round sphere. This is also the intrinsic flat limit. The Gromov-Hausdorff limit, however, does not exist since the diameter diverges to infinity.*

Proof. More precisely the metrics g_i are defined by

$$g_i = h_i(r)^2 dr^2 + f_i^2(r) g_{S^2} \text{ for } r \in [0, \pi] \tag{87}$$

where $f_i(r) = \sin(r)$ and

$$h_i(r) = 1 + i^2 \exp\left(\frac{\left(\frac{1}{2i}\right)^2}{\left(r-\pi+\frac{2}{i}\right)\left(r-\pi+\frac{1}{i}\right)} \right) \chi_{\left[\pi-\frac{2}{i}, \pi-\frac{1}{i}\right)}. \tag{88}$$

Observe that on $r^{-1}[0, \pi - 1/j)$ we have $h_i(r) = 1$ for $i \geq 2j$. So g_i converge smoothly away from p_0 to the standard metric on a sphere, g_∞. The metric and settled completions of $(M \setminus \{p_0\}, g_\infty)$ are both the standard sphere.

Observe that

$$\operatorname{diam}_{g_i}(M_j) \geq \int_0^\pi h_i(r)\, dr = \pi - 2/i + L_i + 1/i \tag{89}$$

where L_i is the length of the spline:

$$L_i = \int_{\pi-2/i}^{\pi-1/i} h_i(r)\, dr \tag{90}$$

$$= i \int_0^1 e^{1/(4u(u-1))}\, du = iL \tag{91}$$

Now taking $U_j = r^{-1}([0, \pi - 1/(2j)])$, we see that

$$\operatorname{diam}_{M_i}(U_j) \le \pi \text{ for } i \ge j, \tag{92}$$

$$\operatorname{Vol}(M_i) \le \operatorname{Vol}(S^3, g_0) + \sin(1/(2i))L_i \le V_0 \tag{93}$$

$$\operatorname{Vol}_{g_i}(\partial W_j) \le 4\pi 1 \tag{94}$$

$$\operatorname{Vol}_{g_i}(M \setminus W_j) \le \pi(1/(2j)^2) + \pi \sin(2/j)^2 L_j \le V_j \tag{95}$$

where $\lim_{j\to\infty} V_j = 0 + \lim_{j\to\infty} \sin^{(} 2/j) jL = 0$. By Theorem 1.3 we have the intrinsic flat limit is settled metric completion which is the sphere. This example has no uniform lower bound on Ricci curvature nor a uniform contractibility function so it demonstrates the necessities of these conditions in all of our theorems which require them to prove the Gromov-Hausdorff limit exists and is the metric completion of $(M \setminus S, g_\infty)$. □

3.8. **Unbounded Boundary Volumes.** Here we have examples demonstrating the necessity of the $\operatorname{Vol}(\partial W)$ conditions in our theorems.

Example 3.12. *There are* (M^3, g_j) *all diffeomorphic to the standard sphere which converge smoothly away from a singular set* $S = \{p_0\}$ *to* $(M \setminus S, g_\infty)$ *with the metric*

$$dr^2 + f^2(r) g_{S^2} \text{ where } r \in [0, \pi) \tag{96}$$

such that $f(r) = \sin(r)$ *on* $[0, \pi/2]$ *and* $\operatorname{Vol}_{g_\infty}(M \setminus S)$ *is finite but*

$$\lim_{r\to\pi} f(r) = \infty. \tag{97}$$

The metric completion agrees with the settled completion of (M_∞, d_∞)*, which is not an integral current space because the area of the boundary is infinite. The diameter of this example is clearly* $\le 2\pi$*. This example demonstrates that (3) of Theorem 1.3 is a necessary condition.*

Proof. Let $g_\infty = dr^2 + f^2(r) g_{S^2}$ where $f(r)$ is a smooth increasing function such that:

$$f(r) = \sin(r) \text{ for } r \in [0, \pi/2] \tag{98}$$

and

$$f(r) = (\pi - r)^{-\frac{1}{4}} \text{ for } r \in [\pi/2 + 1/2, \pi). \tag{99}$$

Then we have:

$$\lim_{r\to\pi} f(r) = \infty \tag{100}$$

and

$$\operatorname{Vol}_{g_\infty}(M \setminus S) = \int_0^\pi \omega_2 f^2(r)\, \mathrm{d}r < \infty. \tag{101}$$

Now let $g_i = dr^2 + f_i^2(r) g_{S^2}$ where, $f_i(r) \leq f(r)$ is a smooth function given by:

$$f_i(r) = f(r) \text{ for } r \in [0, \pi - 1/i] \tag{102}$$

and

$$f_i(r) = \sin(r) \text{ for } r \in [\pi - 1/(2i), \pi]. \tag{103}$$

It is easy to see that g_i converges to g_∞ away from the singular point.

Taking $W_j = r^{-1}([0, \pi - 1/j])$, we see that all conditions of Theorem 1.3 are satisfied except that $\mathrm{Vol}_{g_i}(\partial W_j)$ is not bounded. □

Remark 3.13. *The sequence in Example 3.12 also appears to satisfy uniform local contractibility estimates as there is no cusp effect. The Gromov-Hausdorff limit appears to be the one point completion of* $(M \setminus S, g_\infty)$*. The metric completion of* $(M \setminus S, g_\infty)$ *includes infinitely many new points. So this example may well also prove necessity of boundary volume estimates in Theorem 6.6.*

Remark 3.14. *It is not immediately clear how one may construct similar examples with nonnegative Ricci curvature proving the necessity of (3). A doubly warped product will not suffice where the* W_j *are defined by distance functions will not work because we can control the areas of the level sets of such functions.*

3.9. **Torus to Square.**

Example 3.15. *There are* (M^2, g_j) *all isometric to the flat torus,* $S^1 \times S^1$ *which converge smoothly away from a singular set*

$$S = \left(S^1 \times \{0\}\right) \cup \left(\{0\} \times S^1\right) \subset S^1 \times S^1 \tag{104}$$

to

$$(M \setminus S, g_\infty) = \left((0, 2\pi) \times (0, 2\pi), dt^2 + ds^2\right). \tag{105}$$

So the metric completion and the settled completions are both

$$M_\infty = [0, 2\pi] \times [0, 2\pi] \tag{106}$$

with the standard flat metric, while the intrinsic flat and Gromov Hausdorff limits are the flat torus $S^1 \times S^1$*. Thus the codimension condition and the uniform embeddedness conditions are necessary in all our theorems.*

Proof. Let $W_k = (1/k, 2\pi - 1/k) \times (1/k, 2\pi - 1/k)$. Then, for j large enough:

$$\lambda_{i,j,k} = \sup_{x,y \in W_j} |d_{W_k}(x, y) - d_M(x, y)| = 2\pi - 4/j \tag{107}$$

Therefore,

$$\limsup_{j\to\infty} \limsup_{k\to\infty} \limsup_{i\to\infty} \lambda_{i,j,k} = 2\pi \tag{108}$$

and

$$\limsup_{j\to\infty}\limsup_{i\to\infty}\limsup_{k\to\infty}\lambda_{i,j,k} = 2\pi \tag{109}$$

So we fail uniform embeddedness as well as the codimension 2 condition.

Observe that the sequence satisfies Ricci curvature, contractibility, diameter and volume conditions on M_i because all the M_i are the standard flat torus. Furthermore $\mathrm{Vol}_{g_i}(M\setminus W_j) \le 4/j$ and $\mathrm{Vol}_{g_i}(\partial W_j) \le 4$. □

4. Explicit Estimates with Isometric Embeddings

In this section we construct isometric embeddings of Riemannian manifolds into metric spaces to provide explicit bounds on the Gromov-Hausdorff and intrinsic flat distances between them.

Recall the definition of isometric embedding given in Subsection 2.1. In fact we construct more general mappings.

Definition 4.1. *Let $D > 0$ and M, M' are geodesic metric spaces. We say that $\varphi : M \to M'$ is a D-geodesic embedding if for any smooth minimal geodesic, $\gamma : [0,1] \to M$, of length $\le D$ we have*

$$d_{M'}(\varphi(\gamma(0)), \varphi(\gamma(1))) = L(\gamma). \tag{110}$$

When $D = \mathrm{diam}(M)$, then D-geodesic embeddings are isometric embeddings . The advantage of this more general notion is that it can be applied when M is not complete. This will be essential to proving Theorem 4.6.

4.1. Hemispherical Embeddings.

In this subsection we prove the following key proposition:

Proposition 4.2. *Given a manifold M with Riemannian metrics g_1 and g_2 and $D_1, D_2, t_1, t_2 > 0$. Let $M' = M \times [t_1, t_2]$ and let $\varphi_i : M_i \to M'$ be defined by $\varphi_i(p) = (p, t_i)$. If the metric g' on M' satisfies*

$$g' \ge dt^2 + \cos^2((t-t_i)\pi/D_i)g_i \text{ for } |t-t_i| < D_i/2 \tag{111}$$

and

$$g' = dt^2 + g_i \text{ on } M\times\{t_i\} \subset M' \tag{112}$$

then any geodesic, $\gamma : [0,1] \to M_i$, of length $\le D_i$ satisfies (110). If, the diameter is bounded, $\mathrm{diam}_{g_i}(M) \le D_i$, then φ_i is an isometric embedding.

Furthermore, for $q_1, q_2 \in M$, we have

$$d_{M'}(\varphi_1(q_1), \varphi_2(q_2)) \ge d_{M_i}(q_1, q_2). \tag{113}$$



FIGURE 5. First we see a pair of flat tori, $M_i = (M, g_i)$, isometrically embedded in their own hemispherical suspensions. Then they both isometrically embed into a common M'.

Example 4.3. *Let* $M = S^1 \times [0,1]$ *and let* $g_i = a_i^2 d\theta^2 + b_i^2 dl^2$, *where* $a_1 > a_2 > 0$ *and* $b_2 > b_1 > 0$. *Take*

$$D_i = \mathrm{diam}_{g_i}(M_i) = \sqrt{(\pi a_i)^2 + b_i^2}. \tag{114}$$

By Proposition 4.2, we know that if we can find $t_i \in \mathbb{R}$ *and functions* $a, b : [t_1, t_2] \to \mathbb{R}$ *such that*

$$a(t) \geq \max_{i=1,2} a_i h_i(t) \tag{115}$$

$$b(t) \geq \max_{i=1,2} b_i h_i(t) \tag{116}$$

where $h_i(t) = \max\{\cos((t - t_i)\pi/D_i), 0\}$ *and*

$$a(t_1) = a_1,\ \ a(t_2) = a_2,\ \ b(t_1) = b_1 \ \ \text{and}\ b(t_2) = b_2, \tag{117}$$

then we have isometric embeddings $\varphi_i : (M, g_i) \to (M', g')$ *where*

$$g' = dt^2 + a^2(t)d\theta^2 + b^2(t)dl^2. \tag{118}$$

To obtain (117), we must choose $t_1 - t_2$ *sufficiently large that*

$$a_1 h_1(t_2 - t_1) \leq a_2 \text{ and } b_2 h_2(t_2 - t_1) \leq b_1. \tag{119}$$

Since $a_2/a_1, b_2/b_1 < 1$ *this is achieved by taking*

$$|t_2 - t_1| \geq \max\left\{\frac{D_1}{\pi}\arccos\left(\frac{a_2}{a_1}\right), \frac{D_2}{\pi}\arccos\left(\frac{b_1}{b_2}\right)\right\}. \tag{120}$$

See Figure 5.

Before we prove the proposition we prove the following lemma. Recall that equators in spheres isometrically embed into the hemispheres. Here we create standard isometric embeddings of Riemannian manifolds into hemispherically warped product spaces. The idea comes from Gromov's notions in filling Riemannian manifolds [Gro83].

Lemma 4.4. *Given a Riemannian manifold* (M, g) *and* $D > 0$*. Let* $M' = M \times [0, D/2]$ *and let* $\varphi : M \to M'$ *be defined by* $\varphi(p) = (p, 0)$*. If the metric* g' *on* M' *satisfies*

$$g' \geq dt^2 + \cos^2(t\pi/D)g \text{ on } M' \tag{121}$$

and

$$g' = dt^2 + g \text{ on } M \times \{0\} \subset M' \tag{122}$$

then any geodesic, $\gamma : [0, 1] \to M$*, of length* $\leq D$ *satisfies (110). If* diam$(M) \leq D$ *then* φ *is an isometric embedding.*

Here the hemispherical suspension, M', is a well defined metric space but not necessarily a smooth manifold as can be seen, for example, on the left side of Figure 6. The inspiration for using a hemispherical suspension comes from Gromov's work on filling Riemannian manifolds [Gro83].

Proof. Assume not. There exists a geodesic $\gamma : [0, 1] \to M$ of length $\leq D$, and a curve $\eta : [0, 1] \to M'$ running from $\gamma(0)$ to $\gamma(1)$ of length $L_{g'}(\sigma) < L_g(\gamma)$. If we replace the metric g' by $g'' = dt^2 + \cos^2(t\pi/D)g$, then $L_{g''}(\sigma) < L_g(\gamma)$.

So there exists a curve $C(s) = (x(s), t(s)) \in M \times [0, D/2]$ which is minimizing with respect to g'' between its endpoints $C(0) = (x(0), 0) = \gamma(0)$ and $C(1) = (x(1), 0) = \gamma(1)$ such that

$$L_{g''}(C) \leq L_{g''}(\sigma) \leq L_{g'}(\sigma) < L_g(\gamma) = d_M(x(0), x(1)) \leq D. \tag{123}$$

Since $C : [0, 1] \to M'$ is a minimizing geodesic in a warped product, $x : [0, 1] \to M$ is a minimizing geodesic in M. We choose the parameter s so that x is parametrized proportional to arclength and let h be the length of the geodesic x, so we have $h = d_M(x(0), x(1)) \leq D$. See Figure 6.



Figure 6. On the left we have a smooth torus, M^m, which is warped with a cosine function to create the curved manifold, $M^m \times [0, D/2]$, in which C lies. On the right we see the set U viewed as a subset of a hemisphere created using the same warping function.

Observe that $F : [0,h] \times [0, D/2]$ defined by $F(s,t) = (x(s/h), t)$ is an isometric embedding of a region, U, in the standard round sphere, S^2, of diameter D into M'. That is the metric on U is

$$dt^2 + \cos^2(t\pi/D) ds^2 \tag{124}$$

and for any curve $\gamma : [a,b] \to U$ where $\gamma(u) = (\gamma_s(u), \gamma_t(u))$ we have, by (121),

$$\begin{aligned} g''((F\circ\gamma)'(u),(F\circ\gamma)'(u)) &= (dt^2 + \cos^2(t\pi/D))g(F\circ\gamma)'(u), F\circ\gamma)'(u)) \\ &= \gamma_t'(u)^2 + \cos^2(t\pi/D)g((x\circ\gamma_s)'(u/h),(x\circ\gamma_s)'(u/h))/h^2 \\ &= \gamma_t'(u)^2 + \cos^2(t\pi/D)|\gamma_s'(u)|^2 g(x'(u/h),x'(u/h))/h^2 \\ &= \gamma_t'(u)^2 + \cos^2(t\pi/D)|\gamma_s'(u)|^2 \text{ because } g(x',x') = h^2 \\ &= (dt^2 + \cos^2(t\pi/D)ds^2)(\gamma'(u),\gamma'(u)). \end{aligned}$$

In particular, $L(F\circ x) = L(x)$ since, by (122).

Furthermore $C(s) \subset F(U)$. So $F^{-1}\circ C$ is a curve in S^2 running between $F^{-1}(C(0))$ and $F^{-1}(C(1))$. Thus

$$L_{g''}(C) \ge d_{S^2}(F^{-1}(C(0)), F^{-1}(C(1))) = L(F^{-1}\circ x) = d_M(x(0), x(1)) \tag{125}$$

because $F^{-1}\circ x$ runs along a great circle in S^2 and has length $< D$. This contradicts (123). □

We may now apply Lemma 4.4 to prove Proposition 4.2:

Proof. By Lemma 4.4, (111) and (112), we see that any geodesic, $\gamma : [0,1] \to M_i$, of length $\le D_i$ satisfies (110).

Given $q_1, q_2 \in M$, let $\gamma : [0,1] \to M'$ be a length minimizing geodesic from $\varphi_1(q_1)$ to $\varphi_2(q_2)$. So

$$\gamma(s) = (c(s), t(s)) \in M \times [t_1, t_2]. \tag{126}$$

For $f_i(t) = \cos((t - t_i)\pi/D_i)$

$$\begin{aligned} L_{g'}(\gamma) &= \int_0^1 g'(\gamma'(s),\gamma'(s))\,ds \\ &\ge \int_0^1 \sqrt{t'(s)^2 + \max_{i=1,2} f_i^2(t(s))g_i(c'(s),c'(s))}\,ds \\ &\ge \int_0^1 \sqrt{t'(s)^2 + f_1^2(t(s))g_1(c'(s),c'(s))}\,ds \\ &\ge \int_0^1 \sqrt{t'(s)^2 + f_1^2(t(s))g_1(\bar{c}'(s),\bar{c}'(s))}\,ds \end{aligned}$$

where $\bar{c}$ is a length minimizing geodesic in (M, g_1) from $c(0)$ to $c(1)$ parametrized proportional to arclength of length $h = d_{g_1}(c(0), c(1))$. Thus

$$L_{g'}(\gamma) \geq \int_0^1 \sqrt{t'(s)^2 + f_1^2(t(s))h^2}\, ds. \tag{127}$$

This integral is the length of a curve in a hemisphere of diameter D_1 running from a point $(0, h)$ on the equator to a point $(|t_1 - t_2|, 0)$. So it is greater than or equal to the length of the third side of a triangle opposite a right angle with legs of length $d_{M_1}(q_1, q_2)$ and $|t_1 - t_2|$. Applying the Spherical Law of Cosines rescaled we obtain

$$\begin{aligned} d_{M'}(\varphi_1(q_1), \varphi_2(q_2)) &\geq \frac{D_1}{\pi} \arccos\left(\cos\left(\frac{\pi d_{M_1}(q_1, q_2)}{D_1}\right)\cos\left(\frac{\pi|t_1 - t_2|}{D_1}\right)\right) && (128)\\ &\geq d_{M_1}(q_1, q_2). && (129) \end{aligned}$$

□

4.2. **Estimating Distances between Manifolds.** The Gromov-Hausdorff distance between a pair of metric spaces was estimated in terms of the Lipschitz distance between them in [Gro99]. In [SW11], the intrinsic flat distance distance between a pair of integral current spaces was estimated in terms of the Lipschitz distance between them. Here we give a simple proof estimating these distances between Riemannian manifolds using explicit isometric embeddings into a common Riemannian manifold. Recall Definitions 2.7 and 2.9.

Lemma 4.5. *Suppose $M_1 = (M, g_1)$ and $M_2 = (M, g_2)$ are diffeomorphic oriented precompact Riemannian manifolds and suppose there exists $\epsilon > 0$ such that*

$$g_1(V, V) < (1 + \epsilon)^2 g_2(V, V) \text{ and } g_2(V, V) < (1 + \epsilon)^2 g_1(V, V) \qquad \forall\, V \in TM. \tag{130}$$

Then for any

$$a_1 > \frac{\arccos(1 + \epsilon)^{-1}}{\pi} \operatorname{diam}(M_2) \tag{131}$$

and

$$a_2 > \frac{\arccos(1 + \epsilon)^{-1}}{\pi} \operatorname{diam}(M_1), \tag{132}$$

there is a pair of isometric embeddings $\varphi_i : M_i \to M' = \bar{M} \times [t_1, t_2]$ with a metric as in Proposition 4.2 where $t_2 - t_1 \geq \max\{a_1, a_2\}$.

Thus the Gromov-Hausdorff distance between the metric completions is bounded,

$$d_{GH}(\bar{M}_1, \bar{M}_2) \leq a := \max\{a_1, a_2\}, \tag{133}$$

and the intrinsic flat and scalable intrinsic flat distances between the settled completions are bounded,

$$d_{\mathcal{F}}(M_1', M_2') \leq a\,(V_1 + V_2 + A_1 + A_2), \tag{134}$$

$$d_{s\mathcal{F}}(M_1', M_2') \le (a(V_1 + V_2))^{1/(m+1)} + (a(A_1 + A_2))^{1/(m)} \tag{135}$$

where $V_i = \mathrm{Vol}_m(M_i)$ *and* $A_i = \mathrm{Vol}_{m+1}(\partial M_i)$*.*

Proof. By our choice of a_i we have

$$g_1(V,V) > \cos^2(a_1\pi/D_2) g_2(V,V) \qquad \forall V \in TM \tag{136}$$

and

$$g_2(V,V) > \cos^2(a_2\pi/D_1) g_1(V,V) \qquad \forall V \in TM. \tag{137}$$

Applying Proposition 4.2 and setting $t_1 = 0$ and $t_2 = a$, we have isometric embeddings $\varphi_i : (M, g_i) \to (M', g')$ for any g' satisfying (111) and (112). In fact, the metric g' on M' can be chosen so that

$$g'(V,V) \le dt^2(V,V) + g_1(V,V) + g_2(V,V) \qquad \forall\, V \in TM'. \tag{138}$$

By Definition 2.5 we have,

$$d_{GH}(\bar{M}_1, \bar{M}_2) \le d_H^{M'}(\varphi_1(M_1), \varphi_2(M_2)). \tag{139}$$

For all $r > a$, $\varphi_1(M_1) \subset T_r(\varphi_2(M_2))$ and $\varphi_2(M_2) \subset T_r(\varphi_1(M_1))$, because for all $p \in M$ we have

$$d_{M'}(\varphi_1(p), \varphi_2(p)) = |t_2 - t_1| = a. \tag{140}$$

By Definition 2.4 we have (133).

Recall that to estimate the Intrinsic flat Distance and scalable intrinsic flat Distance we must estimate volumes of a filling manifold and an excess boundary as in (21) and (30). Taking ν to be the unit inward normal to $\partial M' \setminus (M_1 \cup M_2)$ and applying the estimate on g' given in (138) we have

$$\begin{aligned} d_{\mathcal{F}}(M_1', M_2') &\le \mathrm{Vol}(M') + \mathrm{Vol}(\partial M' \setminus (M_1 \cup M_2)) \\ &= \int_{t_1}^{t_2} \int_M \mu_{g'}\, dt + \int_{t_1}^{t_2} \int_{\partial M} \nu \lrcorner \mu_{g'}\, dt \\ &\le |t_2 - t_1|(\mathrm{Vol}(M_1) + \mathrm{Vol}(M_2)) + |t_2 - t_1|(\mathrm{Vol}(\partial M_1) + \mathrm{Vol}(\partial M_2)). \end{aligned}$$

and

$$\begin{aligned} d_{s\mathcal{F}}(M_1', M_2') &\le (\mathrm{Vol}(M'))^{1/(m+1)} + (\mathrm{Vol}(\partial M' \setminus (M_1 \cup M_2)))^{1/m} \\ &= \left(\int_{t_1}^{t_2} \int_M \mu_{g'}\, dt\right)^{1/(m+1)} + \left(\int_{t_1}^{t_2} \int_{\partial M} \nu \lrcorner \mu_{g'}\, dt\right)^{1/m} \\ &\le (|t_2 - t_1|(\mathrm{Vol}(M_1) + \mathrm{Vol}(M_2)))^{1/(m+1)} \\ &\qquad + (|t_2 - t_1|(\mathrm{Vol}(\partial M_1) + \mathrm{Vol}(\partial M_2)))^{1/m}. \end{aligned}$$

□

4.3. **Appending Regions without Smooth Approximations.** Now we examine pairs of precompact oriented manifolds (M_1, g_1) and (M_2, g_2) which are not diffeomorphic but have diffeomorphic regions $U_i \subset M_i$. That is, there is a common smooth manifold with boundary U and diffeomorphisms $\psi_i : U \to U_i \subset M_i$. Then we may apply Proposition 4.2 and Lemma 4.5 to the regions U_i to estimate the distances between the metric and settled completions of the M_i. Recall also Definitions 2.7 and 2.9. Recall also the distinction between the intrinsic length metric, d_U, and the restricted metric d_M, on a region $U \subset M$ and the corresponding diameters, $\operatorname{diam}_M(U) \le \operatorname{diam}_U(U)$, in Definition 2.1.

Theorem 4.6. *Suppose $M_1 = (M, g_1)$ and $M_2 = (M, g_2)$ are oriented precompact Riemannian manifolds with diffeomorphic subregions $U_i \subset M_i$ and diffeomorphisms $\psi_i : U \to U_i$ such that*

$$\psi_1^* g_1(V, V) < (1+\epsilon)^2 \psi_2^* g_2(V, V) \qquad \forall\, V \in TU \tag{141}$$

and

$$\psi_2^* g_2(V, V) < (1+\epsilon)^2 \psi_1^* g_1(V, V) \qquad \forall\, V \in TU. \tag{142}$$

Taking the extrinsic diameters,

$$D_{U_i} = \sup\{\operatorname{diam}_{M_i}(W) : \textit{ W is a connected component of } U_i\} \le \operatorname{diam}(M_i), \tag{143}$$

we define a hemispherical width,

$$a > \frac{\arccos(1+\epsilon)^{-1}}{\pi} \max\{D_{U_1}, D_{U_2}\}. \tag{144}$$

Taking the difference in distances with respect to the outside manifolds,

$$\lambda = \sup_{x,y \in U} |d_{M_1}(\psi_1(x), \psi_1(y)) - d_{M_2}(\psi_2(x), \psi_2(y))|, \tag{145}$$

we define heights,

$$h = \sqrt{\lambda(\max\{D_{U_1}, D_{U_2}\} + \lambda/4)} \tag{146}$$

and

$$\bar{h} = \max\{h, \sqrt{\epsilon^2 + 2\epsilon}\, D_{U_1}, \sqrt{\epsilon^2 + 2\epsilon}\, D_{U_2}\}. \tag{147}$$

Then the Gromov-Hausdorff distance between the metric completions is bounded,

$$d_{GH}(\bar{M}_1, \bar{M}_2) \le a + 2\bar{h} + \max\left\{d_H^{M_1}(U_1, M_1), d_H^{M_2}(U_2, M_2)\right\} \tag{148}$$

and the intrinsic flat distance between the settled completions is bounded,

$$\begin{aligned} d_{\mathcal{F}}(M_1', M_2') \quad \le \quad & \left(2\bar{h} + a\right)\left(\operatorname{Vol}_m(U_1) + \operatorname{Vol}_m(U_2) + \operatorname{Vol}_{m-1}(\partial U_1) + \operatorname{Vol}_{m-1}(\partial U_2)\right) \\ & + \operatorname{Vol}_m(M_1 \setminus U_1) + \operatorname{Vol}_m(M_2 \setminus U_2). \end{aligned}$$

and the scalable intrinsic flat distance is bounded,

$$\begin{aligned} d_{s\mathcal{F}}(M_1', M_2') &\le \Big((\mathrm{Vol}_m(U_1) + \mathrm{Vol}_m(U_2)) \big(\bar{h} + a\big) \Big)^{1/(m+1)} \\ &\quad + \Big(\big(2\bar{h} + a\big) (\mathrm{Vol}_{m-1}(\partial U_1) + \mathrm{Vol}_{m-1}(\partial U_2)) \\ &\quad + \mathrm{Vol}_m(M_1 \setminus U_1) + \mathrm{Vol}_m(M_2 \setminus U_2) \Big)^{1/m}. \end{aligned}$$

Figure 2 may be viewed as an application of this theorem. It should be noted that this theorem is an improvement on the Bridge Method Lemma A.2 of [SW11] in two respects. First, we allow U_1 and U_2 not isometric, and secondly we loosen the diameter bounds of that method asking only for control on the λ defined here.

Recall in Definition 2.1, that two different metrics are defined on a connected subdomain, $U \subset M$. When U is is also convex, these two metrics agree. Theorem 4.6 does not require the subdomains to be connected or convex, and so the proof becomes quite difficult. Before we prove this theorem we state and prove a special case with stronger estimates.

Theorem 4.7. *Suppose $M_1 = (M, g_1)$ and $M_2 = (M, g_2)$ are oriented Riemannian manifolds with diffeomorphic convex subregions $U_i \subset M_i$ and diffeomorphisms $\psi_i : U \to U_i$ such that*

$$\psi_1^* g_1(V, V) < (1+\epsilon)^2 \psi_2^* g_2(V, V) \qquad \forall\, V \in TU \tag{149}$$

and

$$\psi_2^* g_2(V, V) < (1+\epsilon)^2 \psi_1^* g_1(V, V) \qquad \forall\, V \in TU. \tag{150}$$

Then for any

$$a_1 > \frac{\arccos(1+\epsilon)^{-1}}{\pi} \operatorname{diam}_{U_2}(U_2) \tag{151}$$

and

$$a_2 > \frac{\arccos(1+\epsilon)^{-1}}{\pi} \operatorname{diam}_{U_1}(U_1), \tag{152}$$

there is a pair of isometric embeddings $\varphi_i : U_i \to M'$ where $M' = U \times [t_1, t_2]$ where $t_2 - t_1 = \max\{a_1, a_2\}$ such that $\varphi_i(x) = (x, t_i)$. Furthermore, these isometric embeddings extend to isometric embeddings $\varphi : M_i \to Z'$, where Z' is a length metric space defined by gluing M_i to M' along U_i.

In particular the Gromov-Hausdorff distance between the metric completions is bounded,

$$d_{GH}(\bar{M}_1, \bar{M}_2) \le \max\{a_1, a_2\} + \max\left\{ d_H^{M_1}(U_1, M_1), d_H^{M_2}(U_2, M_2) \right\} \tag{153}$$

and the intrinsic flat distance between the settled completions is bounded,

$$\begin{aligned} d_{\mathcal{F}}(M_1', M_2') &\le \max\{a_1, a_2\} (\mathrm{Vol}(U_1) + \mathrm{Vol}(U_2) + \mathrm{Vol}(\partial U_1) + \mathrm{Vol}(\partial U_2)) \\ &\quad + \mathrm{Vol}(M_1 \setminus U_1) + \mathrm{Vol}(M_2 \setminus U_2), \end{aligned}$$

and the scalable intrinsic flat distance is bounded,

$$\begin{aligned} d_{s\mathcal{F}}(M_1', M_2') \;\le\; & \left(\max\{a_1, a_2\} \left(\mathrm{Vol}(U_1) + \mathrm{Vol}(U_2)\right) \right)^{1/(m+1)} \\ & + \left(\max\{a_1, a_2\} \left(\mathrm{Vol}(\partial U_1) + \mathrm{Vol}(\partial U_2)\right) + \mathrm{Vol}(M_1 \setminus U_1) + \mathrm{Vol}(M_2 \setminus U_2) \right)^{1/(m)}. \end{aligned}$$

Proof. The metric g' on M' is defined by applying Proposition 4.2 and Lemma 4.5 to the diffeomorphic regions, U_1 and U_2, using the D_i as defined above $\varphi_i : U_i \to M'$ are isometric embeddings. We can choose g' satisfying (138).

We must verify that the M_i isometrically embed into Z' constructed as in the statement of the theorem. To see this we take any $x, y \in M_1$ and a shortest curve $C \subset Z'$ running between $\varphi_1(x)$ and $\varphi_1(y)$. If the curve never enters $\varphi_2(M_2 \setminus U_2)$ then $d_{M_1}(x, y) = d_{Z'}(\varphi_1(x), \varphi_1(y))$ by Lemma 4.5 and Lemma A.1 in the Appendix of [SW11] applied to $\varphi_1(M_1) \cup M' \subset Z'$. If the curve does enter $\varphi_2(M_2 \setminus U_2)$ then we have a length minimizing curve which leaves $\bar{U}_2$ contradicting the fact that it is convex. The same argument may be repeated to prove $\varphi_2 : M_2 \to Z'$ is an isometric embedding.

So now we may estimate the Gromov-Hausdorff distance as in Remark 2.6. Let $r_i = d_H^{M_i}(M_i, U_i)$. We claim

$$d_{GH}(M_1, M_2) \le d_H^{Z'}(\varphi_1(M_1), \varphi_2(M_2)) \le |t_1 - t_2| + \max\{r_i\}. \tag{154}$$

Fix any $\delta > 0$. Then any point $p \in M_1$ has a point $q \in U_1$ such that $d(p, q) \le r_1 + \delta$. Furthermore, $\varphi_1(q) = (q, t_1) \in M' \subset Z'$ so

$$d_{Z'}(\varphi_1(q), (q, t_2)) = |t_2 - t_1| \tag{155}$$

and $(q, t_2) \subset \varphi_2(U_2) \subset \varphi_2(M_2)$. Thus

$$\varphi_1(M_1) \subset T_{r_1+\delta+|t_2-t_1|}(\varphi(M_2)) \tag{156}$$

and similarly

$$\varphi_2(M_2) \subset T_{r_2+\delta+|t_2-t_1|}(\varphi(M_1)). \tag{157}$$

The claim follows by taking $\delta \to 0$.

We bound the intrinsic flat distance as in Remark 2.8 taking M' to be the filling manifold with the metric g' defined in Lemma 4.5 satisfying (138). We apply the same estimates as in Lemma 4.5 to bound the volumes of these regions, only now we add in the additional volume terms coming from the additional components of the excess boundary $M_i \setminus U_i$.

We bound the scalable intrinsic flat distance as in Remark 2.11. Again we include the additional components of the excess boundary but insert them into the summand with an exponent of $1/m$ since these are m dimensional boundary regions and the scalable flat distance is 1 dimensional. □

We now prove Theorem 4.6. To prove this theorem we adapt the proof of the convex case and the proof of Lemma A.2 in [SW11]. It is essential to possibly push the two manifolds further apart than required simply to isometrically embed the U_i into M' as a short cut for a path between points in $\varphi_1(M_1)$ might be found within $\varphi_2(M_2 \setminus U_2)$.

Proof. For each corresponding pair of connected components $U_{\alpha,i}$ of U_i, we create a hemispherically defined filling bridge M'_α diffeomorphic to $U_{\alpha_i,i} \times [0,a]$ with metric g'_α satisfying (110) by applying Proposition 4.2 and Lemma 4.5 using the $a_i = a_i(\alpha)$ defined there for that particular connected component, U_α and $D_i = D_{U_i}$. Observe that all $a_i \le a$, so we may take $|t_1 - t_2| = a$ for all the connected components. Any minimal geodesic $\gamma : [0,1] \to U_{\alpha,i}$ of length $\le D_{U_i} \le \operatorname{diam}_{M_i}(U_i)$ satisfies (110).

We take the disjoint unions of these bridges to be M'. So it has a metric g' satisfying (138). Observe that the boundary of M' is $(U, g_1) \cup (U, g_2) \cup (\partial U \times [0,a], g')$. So that

$$\begin{aligned}
\operatorname{Vol}_m(M') &= \sum_\alpha \operatorname{Vol}_m(M'_\alpha) && (158)\\
&\le \sum_\alpha a(\operatorname{Vol}_m(U_{\alpha,1}) + \operatorname{Vol}_m(U_{\alpha,2})) && (159)\\
&\le a(\operatorname{Vol}_m(U_1) + \operatorname{Vol}_m(U_2)) && (160)
\end{aligned}$$

and

$$\operatorname{Vol}_m\left(\partial M' \setminus (\varphi_1(U_1) \cup \varphi_2(U_2)\right) \le a\left(\operatorname{Vol}_{m-1}(\partial U_1) + \operatorname{Vol}_{m-1}(\partial U_2)\right) \tag{161}$$

as in Lemma 4.5.

We cannot directly glue M_i to M' and obtain an isometric embedding because our regions are not convex. On either end of the filling bridges, we glue isometric products $U_\alpha \times [0,\bar{h}]$ with metric $dt^2 + g_i$, so that all the bridges are extended by an equal length on either side. This creates a Lipschitz manifold,

$$M'' = (U_1 \times [0,\bar{h}]) \sqcup_{U_1} M' \sqcup_{U_2} (U_2 \times [0,\bar{h}]). \tag{162}$$

We then define $\varphi_i : U_i \to M$ such that

$$\begin{aligned}
\varphi_1(x) &= (x,0) \in U_1 \times [0,\bar{h}] && (163)\\
\varphi_2(x) &= (x,\bar{h}) \in U_2 \times [0,\bar{h}] && (164)
\end{aligned}$$

as in Figure 7. Then by (158) and (161), we have

$$\begin{aligned}
\operatorname{Vol}_{m+1}(M'') &= \operatorname{Vol}_{m+1}(M') + \bar{h}(\operatorname{Vol}_m(U_1) + \operatorname{Vol}_m(U_2)) && (165)\\
&\le (a+\bar{h})(\operatorname{Vol}_m(U_1) + \operatorname{Vol}_m(U_2)) && (166)
\end{aligned}$$

and $\operatorname{Vol}_m\left(\partial M'' \setminus (\varphi_1(U_1) \cup \varphi_2(U_2)\right) =$

$$\begin{aligned}
&= \operatorname{Vol}_m\left(\partial M' \setminus (\varphi_1(U_1) \cup \varphi_2(U_2)\right) + \bar{h}(\operatorname{Vol}_{m-1}(\partial U_1) + \operatorname{Vol}_{m-1}(\partial U_2)) && (167)\\
&\le (a+\bar{h})(\operatorname{Vol}_{m-1}(\partial U_1) + \operatorname{Vol}_{m-1}(\partial U_2)). && (168)
\end{aligned}$$

Finally we glue M_1 and M_2 to the far ends of M'' along $\varphi_i(U_i)$ to create a connected length space

$$Z = \bar{M}_1 \sqcup_{U_1} M'' \sqcup_{U_2} \bar{M}_2 \tag{169}$$

where distances in Z are defined by taking the infimum of lengths of curves as usual. See Figure 7. We will refer to each connected component, M''_α of M'' as the filling bridge corresponding to U_α.



FIGURE 7. Creating Z for Theorem 4.6.

We must prove that $\varphi_1 : M_1 \to Z$ mapping M_1 into its copy in Z is an isometric embedding. To see this we take any $x, y \in M_1$ and a shortest curve $C \subset Z$ running between $\varphi_1(x)$ and $\varphi_1(y)$. As in the convex proof, our only concern is the possibility that C passes into $\varphi_2(M_2 \setminus U_2)$.

If the minimizing curve never crosses a filling bridge, then we claim it has the same length as a curve in $\varphi_1(M_1)$. To see this, we take any $s_2 > s_1 \in [0, 1]$ such that $C(s_1), C(s_2) \in \varphi_1(M_1)$ and $C((s_1, s_2)) \subset Z \setminus \varphi_1(M_1)$. Since C is assumed not to cross a bridge (not to enter $\varphi_2(M_2)$, then $C(s_1) = \varphi(x_1)$ and $C(s_2) = \varphi(x_2)$ where x_1, x_2 lie in the same connected component, $U_{\alpha,1}$, of U_1. Since $C([s_1, s_2])$ is a minimizing curve it has length

$$\leq d_{M''}(C(s_1), C(s_2)) \leq d_{M_1}(x_1, x_2) \leq \operatorname{diam}_{M_1}(U_{\alpha,1}) \leq D_{U_1}. \tag{170}$$

By (110), a minimal geodesic from x_1 to x_2 lying in $U_{\alpha,1}$ has the same length as $C([s_1, s_2])$. So we may replace this segment of C with the image of this minimal geodesic.

On the other hand, if the minimizing curve crosses a filling bridge all the way to $\varphi_2(M_2)$, then we may carefully apply the choice of $\bar{h}$ to reach a contradiction as in the left hand side of Figure 8. We define the following points $0 \leq t_1 < t_2 \leq t_3 <$

$t_4 \le 1$ such that

$$t_1 = \inf\{t : \ C(t) \in Cl(Z \setminus \varphi_1(M_1))\}, \tag{171}$$

$$t_2 = \min\{t : \ C(t) \in \varphi_2(M_2)\}, \tag{172}$$

$$t_4 = \min\{t > t_2 : \ C(t) \in \varphi_1(M_1)\}, \tag{173}$$

$$t_3 = \max\{t \in [t_2, t_4) : \ C(t) \in \varphi_2(M_2)\} \tag{174}$$

so that $C([t_1, t_2])$ and $C([t_3, t_4])$ are geodesic segments lying within filling bridges:

$$C([t_1, t_2]) \subset M''_{\alpha_{1,2}} \qquad C([t_3, t_4]) \subset M''_{\alpha_{3,4}}. \tag{175}$$



FIGURE 8. Why length minimizing curves cannot cross bridges in the proof of Theorem 4.6.

Observe that there are points $p_1, p_4 \in U$ and $p_2, p_3 \in \partial(U)$ such that

$$\varphi_1(\psi_1(p_1)) = C(t_1) \qquad \varphi_1(\psi_1(p_4)) = C(t_4) \tag{176}$$

$$\varphi_2(\psi_2(p_2)) = C(t_2) \qquad \varphi_2(\psi_2(p_3)) = C(t_3). \tag{177}$$

Observe that since $C([t_2, t_3]) \subset \varphi_2(M_2)$ we know the length of this segment is

$$d_Z(C(t_2), C(t_3)) = d_{M_2}(\psi_2(p_2), \psi_2(p_3)) \ge d_{M_1}(\psi_1(p_2), \psi_1(p_3)) - \lambda \tag{178}$$

by the definition of λ in (145).

We claim that the lengths of the other segments are

$$d_{M''}(C(t_1), C(t_2)) > \sqrt{d_{M_1}(p_1, p_2)^2 + h^2} \tag{179}$$

and

$$d_{M''}(C(t_3), C(t_4)) > \sqrt{d_{M_1}(p_3, p_4)^2 + h^2}\,. \tag{180}$$

Once we prove this claim, we see that by the definition of h we have

$$d_{M''}(C(t_1), C(t_2)) > d_{M_1}(p_1, p_2) + \lambda/2, \tag{181}$$

$$d_{M''}(C(t_3), C(t_4)) > d_{M_1}(p_3, p_4) + \lambda/2. \tag{182}$$

This combined with (178) implies that

$$\begin{aligned} L(C([t_1, t_4]) &= d_{M'}(C(t_1), C(t_2)) + d_Z(C(t_2), C(t_3)) + d_{M''}(C(t_3), C(t_4)) \\ &> d_{M_1}(p_1, p_2) + \lambda/2 + d_{M_1}(p_2, p_3) - \lambda + d_{M_1}(p_3, p_4) + \lambda/2 \\ &\geq d_{M_1}(p_1, p_2) + d_{M_1}(p_2, p_3) d_{M_1}(p_3, p_4) \\ &\geq d_{M_1}(p_1, p_4) = d_{\varphi_1(M_1)}(C(t_1), C(t_4)) \\ &\geq d_Z(C(t_1), C(t_4)), \end{aligned}$$

which is a contradicts the fact that C was minimizing.

So we need only prove our claim in (179) and (180) to see that $\varphi_1 : M_1 \to Z$ is an isometric embedding. This claim concerns a minimizing geodesic lying in a single connected component of the filling bridges,

$$\gamma : [0, 1] \to (U_{\alpha,1} \times [0, \bar{h}]) \sqcup_{U_{\alpha,1}} M'_\alpha \sqcup_{U_{\alpha,2}} (U_{\alpha,2} \times [0, \bar{h}]) \tag{183}$$

such that

$$\gamma(0) = (q_0, 0) \in U_{\alpha,1} \times \{0\} \subset \varphi_1(M_1) \tag{184}$$

and

$$\gamma(1) = (q_3, \bar{h}) \in U_{\alpha,2} \times \{\bar{h}\} \subset \varphi_2(M_2). \tag{185}$$

Consult the right hand side of Figure 8. Let $0 < s_1 < s_2 < 1$ be chosen so that

$$\gamma(s_1) = (q_1, \bar{h}) \in U_{\alpha,1} \times \{\bar{h}\} \subset \partial M'_\alpha \tag{186}$$

and

$$\gamma(s_2) = (q_2, 0) \in U_{\alpha,2} \times \{0\} \subset \partial M'_\alpha. \tag{187}$$

Then by (113), we have

$$L_{g'}(\gamma([s_1, s_2]) = d_{g'}(\gamma(s_1), \gamma(s_2)) \geq d_{g_1}(q_1, q_2), \tag{188}$$

so that

$$\begin{aligned}
L(\gamma) &= d_Z(\gamma(0),\gamma(s_1)) + d_Z(\gamma(s_1),\gamma(s_2)) + d_Z(\gamma(s_2),\gamma(1)) \\
&= \sqrt{d_{U_1}(q_0,q_1)^2+\bar{h}^2} + d_{U_1}(q_1,q_2) + \sqrt{d_{U_2}(q_2,q_3)^2+\bar{h}^2} \\
&\geq \sqrt{d_{U_1}(q_0,q_1)^2+h^2} + d_{U_1}(q_1,q_2) \\
&+ \quad \sqrt{d_{U_1}(q_2,q_3)^2/(1+\epsilon)^2+(\epsilon^2+2\epsilon)D_{U_1}^2} \\
&\geq \sqrt{d_{M_1}(q_0,q_1)^2+h^2} + d_{M_1}(q_1,q_2) \\
&+ \quad \sqrt{d_{M_1}(q_2,q_3)^2/(1+\epsilon)^2+(\epsilon^2+2\epsilon)d_{M_1}(q_3,q_4)^2} \\
&\geq \sqrt{d_{M_1}(q_0,q_1)^2+h^2} + d_{M_1}(q_1,q_2) + \sqrt{d_{M_1}(q_2,q_3)^2} \\
&> \sqrt{(d_{M_1}(q_0,q_1)+d_{M_1}(q_1,q_2)+d_{M_1}(q_2,q_3))^2+h^2} \\
&\geq \sqrt{d_{M_1}(q_0,q_3)^2+h^2}.
\end{aligned}$$

This gives us (179) and (180). Thus we have proven $\varphi_1 : M_1 \to Z$ is an isometric embedding and the same follows for $\varphi_2 : M_2 \to Z$.

So now we may estimate the Gromov-Hausdorff distance: Let $r_i = d_H^{M_i}(M_i, U_i)$. We claim

$$d_{GH}(M_1,M_2) \leq d_H^Z(\varphi_1(M_1),\varphi_2(M_2) \leq \bar{h}+\bar{h}+a+\max\{r_i\}\,. \tag{189}$$

Fix any $\delta > 0$. Then any point $p \in M_1$ has a point $q \in U_1$ such that $d(p,q) \leq r_1+\delta$. Furthermore,

$$d_Z(\varphi_1(q),\varphi_2(q)) \leq a+\bar{h}+\bar{h} \tag{190}$$

and $\varphi_2(q) \subset \varphi_2(U_2) \subset \varphi_2(M_2)$. Thus

$$\varphi_1(M_1) \subset T_{r_1+\delta+a+\bar{h}+\bar{h}}(\varphi(M_2)). \tag{191}$$

and similarly

$$\varphi_2(M_2) \subset T_{r_2+\delta+a+\bar{h}+\bar{h}}(\varphi(M_1)). \tag{192}$$

The claim follows by taking $\delta \to 0$.

To bound the intrinsic flat distance and scalable intrinsic flat distance, we take $B^{m+1} = M''$ to be the filling manifold and then the excess boundary is

$$A^m = \varphi_1(M_1 \setminus U_1) \cup \varphi_2(M_2 \setminus U_2) \cup \partial M'' \setminus (\varphi_1(U_1) \cup \varphi_2(U_2)) \tag{193}$$

so that with appropriate orientations we have

$$\int_{\varphi_1(M_1)} \omega - \int_{\varphi_2(M_2)} \omega = \int_{B^{m+1}} d\omega + \int_{A^m} \omega. \tag{194}$$

The volumes of these manifolds have been computed in (167) and (165). So as in Remark 2.8 we have

$$\begin{aligned} d_{\mathcal{F}}(M_1, M_2) &\le \operatorname{Vol}_m(U_1)\left(\bar{h}+a\right) + \operatorname{Vol}_m(U_2)\left(\bar{h}+a\right) \\ &\quad + \left(\bar{h}+a\right)\operatorname{Vol}_{m-1}(\partial U_1) + \left(\bar{h}+a\right)\operatorname{Vol}_{m-1}(\partial U_2) \\ &\quad + \operatorname{Vol}_m(M_1 \setminus U_1) + \operatorname{Vol}_m(M_2 \setminus U_2). \end{aligned}$$

The scalable intrinsic flat distance is bounded as in Remark 2.11 so that we have

$$\begin{aligned} d_{s\mathcal{F}}(M_1, M_2) &\le \left(\operatorname{Vol}_m(U_1)\left(\bar{h}+a\right) + \operatorname{Vol}_m(U_2)\left(\bar{h}+a\right)\right)^{1/(m+1)} \\ &\quad + \Big(\left(\bar{h}+a\right)\operatorname{Vol}_{m-1}(\partial U_1) + \left(\bar{h}+a\right)\operatorname{Vol}_{m-1}(\partial U_2) \\ &\quad + \operatorname{Vol}_m(M_1 \setminus U_1) + \operatorname{Vol}_m(M_2 \setminus U_2)\Big)^{1/m}. \end{aligned}$$

□

5. Intrinsic Flat Limits

In this section we examine sequences of Riemannian manifolds which converge smoothly away from singular sets and their intrinsic flat limits proving Theorem 1.3. This theorem will be shown to be consequences of the following more powerful theorem which requires a condition on the embeddings of the exhaustion in the manifold:

Definition 5.1. *Given a sequence of Riemannian manifolds $M_i = (M, g_i)$ and an open subset, $U \subset M$, a connected precompact exhaustion, W_j, of U satisfying (2) is uniformly well embedded if*

$$\lambda_{i,j,k} = \sup_{x,y\in W_j} |d_{(W_k,g_i)}(x,y) - d_{(M,g_i)}(x,y)| \tag{195}$$

has

$$\limsup_{j\to\infty}\limsup_{k\to\infty}\limsup_{i\to\infty}\lambda_{i,j,k} \le \lambda_0 \tag{196}$$

and

$$\limsup_{k\to\infty}\lambda_{i,j,k} = \lambda_{i,j} \text{ where } \limsup_{i\to\infty}\lambda_{i,j} = \lambda_j \text{ and } \lim_{j\to\infty}\lambda_j = 0. \tag{197}$$

Theorem 5.2. *Let $M_i = (M, g_i)$ be a sequence of Riemannian manifolds such that there is a closed subset, S, and a uniformly well embedded connected precompact exhaustion, W_j, of $M \setminus S$ satisfying (2) such that g_i converge smoothly to g_∞ on each W_j with*

$$\operatorname{diam}_{M_i}(W_j) \le D_0 \qquad \forall i \ge j, \tag{198}$$

$$\operatorname{Vol}_{g_i}(\partial W_j) \le A_0 \tag{199}$$

and

$$\mathrm{Vol}_{g_i}(M \setminus W_j) \le V_j \text{ where } \lim_{j\to\infty} V_j = 0. \tag{200}$$

Then

$$\lim_{j\to\infty} d_{\mathcal{F}}(M'_j, N') = 0 \tag{201}$$

where N' is the settled completion of $(M \setminus S, g_\infty)$.

In the first subsection, we prove a technical proposition demonstrating that the intrinsic flat limit of a connected precompact exhaustion of an open set in a fixed Riemannian manifold to the metric completion of that open set [Proposition 5.4]. This theorem is shown to be false for Gromov-Hausdorff limits [Example 5.5].

The second subsection, we complete the proof of Theorem 5.2 applying Proposition 5.4 and Theorem 4.6.

The third subsection contains a proof of Lemma 5.7 concerning manifolds with singular sets of codimension two. This final lemma combined with Theorem 5.2 proves Theorem 1.3.

Remark 5.3. *In Example 3.4 we see that it is necessary to assume that the exhaustion is connected in Theorem 5.2. The excess volume bound in (4) is shown to be necessary in Example 3.7 and Example 3.8, which has no intrinsic flat limit. The uniform bound on the boundary volumes, (3), is seen to be necessary in Example 3.12. All these examples have codimension 2 singular sets and show the necessity of these hypothesis for Theorem 1.3 as well. The uniform embeddedness hypothesis of Theorem 5.2 and the codimension two condition of Theorem 1.3 are seen to be necessary for their respective theorems in Example 3.15.*

5.1. **Creating Spaces from Exhaustions.**

In this section we examine the construction of the limit space from a sequence of precompact open sets. One may view this section as a technical subsection. Recall that a connected precompact exhaustion of a domain satisfies (2).

Proposition 5.4. *Let W_j be a connected precompact exhaustion of a Riemannian manifold, N, with fixed Riemannian metric, g_N. If we assume that* $\mathrm{diam}(N) \le D_0$, $\mathrm{Vol}(W_j) \le \mathrm{Vol}(N) \le V_0$ *and* $\mathrm{Vol}(\partial W_j) \le A_0$ *then the settled completion $N' \subset \bar{N}$ satisfies*

$$\lim_{j\to\infty} d_F((W'_j, d_{W_j}), (N', d_{N'})) = 0. \tag{202}$$

where d_{W_j} is the induced length metric on W_j defined by the Riemannian metric g_N and W'_j is the settled completion of W_j with respect to d_{W_j}.

The connectedness is essential to this theorem as can be seen in Example 3.4. Interestingly, one does not obtain Gromov-Hausdorff convergence under these conditions. There need not even exist a Gromov-Hausdorff limit of $(\bar{W}_j, d_{W_j})$. See Example 5.5 below.

Proof. We first verify that we can apply Theorem 4.6 with $M_1 = W_k$ and $M_2 = N$ and $U_1 = W_i \subset W_k$ for $i < k$ and $U_2 = W_i \subset N$. Note that $\epsilon = 0$ and the hemispherical width a can be taken to be 0 because U_i have the same Riemannian metric, g_N.

We claim

$$\lim_{j\to\infty} \mathrm{Vol}(N \setminus W_j) = 0. \tag{203}$$

Since N is an open manifold of finite volume

$$\mathrm{Vol}(N) = \sum_{k=1}^{\infty} \mathrm{Vol}(W_k \setminus W_{k-1}), \tag{204}$$

so

$$\lim_{j\to\infty} \sum_{k=j+1}^{\infty} \mathrm{Vol}(W_k \setminus W_{k-1}) = 0. \tag{205}$$

However

$$\mathrm{Vol}(N \setminus W_j) = \sum_{k=j+1}^{\infty} \mathrm{Vol}(W_k \setminus W_{k-1}) \tag{206}$$

so we have our claim.

Let $k > i$ and let

$$\lambda_{i,k} = \sup_{x,y\in W_i} |d_{W_k}(x,y) - d_N(x,y)|. \tag{207}$$

Then

$$D_{U_1} = \mathrm{diam}_{W_k}(W_i) \le \mathrm{diam}_N W_i + \lambda_{i,k} \le D_0 + \lambda_{i,k}. \tag{208}$$

and,

$$D_{U_2} = \mathrm{diam}_N W_i \le D_0. \tag{209}$$

We claim that for fixed i,

$$\lim_{k\to\infty} \lambda_{i,k} = 0. \tag{210}$$

First note that $\lambda_{i,k}$ is decreasing in k because

$$d_{W_k}(x,y) \ge d_{W_{k+1}}(x,y) \ge d_N(x,y). \tag{211}$$

If the limit is not zero in (210) then let

$$\epsilon' = \inf_k \lambda_{i,k} > 0. \tag{212}$$

Since $\bar{W}_i$ is compact, there exists $x_{i,k}, y_{i,k} \subset \bar{W}_i$ achieving the supremum in (207). Taking k to infinity, a subsequence converges to $x_i, y_i \subset \bar{W}_i$ with respect to $d_{\bar{W}_i}$. Let $\gamma_i \subset N$ be a curve from x_i to y_i such that

$$L(\gamma_i) \le d_N(x_i,y_i) + \epsilon'/5. \tag{213}$$

Since W_k exhaust N, there exists $N_{\epsilon'}$ sufficiently large that

$$\gamma_i \subset W_k \qquad \forall k \geq N_{\epsilon'}. \tag{214}$$

Thus

$$d_{W_k}(x_i, y_i) \leq d_N(x_i, y_i) + \epsilon'/5 \qquad \forall k \geq N_{\epsilon'}. \tag{215}$$

Now take k from the subsequence sufficiently large that we have

$$\begin{aligned} d_{\bar{W}_k}(x_{i,k}, x_i) &\leq d_{\bar{W}_i}(x_{i,k}, x_i) < \epsilon'/5 && (216)\\ d_{\bar{W}_k}(y_{i,k}, y_i) &\leq d_{\bar{W}_i}(y_{i,k}, y_i) < \epsilon'/5 && (217) \end{aligned}$$

and then

$$\begin{aligned} d_N(x_{i,k}, y_{i,k}) &\leq d_{W_k}(x_{i,k}, y_{i,k})\\ &\leq d_{W_k}(x_{i_k}, x_i) + d_{W_k}(x_i, y_i) + d_{W_k}(y_i, y_{i_k})\\ &< d_N(x_i, y_i) + 3\epsilon'/5\\ &\leq d_N(x_{i_k}, x_i) + d_N(x_i, y_i) + d_N(y_i, y_{i_k}) + 3\epsilon'/5\\ &\leq d_{W_k}(x_{i_k}, x_i) + d_N(x_i, y_i) + d_{W_k}(y_i, y_{i_k}) + 3\epsilon'/5\\ &\leq d_{W_k}(x_{i_k}, x_i) + d_N(x_i, y_i) + d_{W_k}(y_i, y_{i_k}) + 3\epsilon'/5\\ &< 5\epsilon'/5 = \epsilon' \end{aligned}$$

and $\lambda_{i,k} < 3\epsilon'/5$ which is a contradiction.

By (208), (209) and (210), we know that for fixed i,

$$\lim_{k\to\infty} \bar{h}_{i,k} = 0 \tag{218}$$

where $\bar{h}_{i,k}$ is defined as in (146)-(147) with $\lambda = \lambda_{i,j}$ and $\epsilon = 0$.

By Theorem 4.6, the intrinsic flat distance is bounded,

$$\begin{aligned} d_{\mathcal{F}}(N', W_j) &\leq \left(\bar{h}_{i,j}\right)(2\,\mathrm{Vol}(W_i) + 2\,\mathrm{Vol}(\partial W_i)) + \mathrm{Vol}(N \setminus W_i) + \mathrm{Vol}(W_j \setminus W_i)\\ &\leq \left(\bar{h}_{i,j}\right)(2V_0 + 2A_0) + \mathrm{Vol}(N \setminus W_i) + \mathrm{Vol}(N \setminus W_i) \end{aligned}$$

By (203), for any ϵ" > 0 there exists i sufficiently large that

$$d_{\mathcal{F}}(N', W_j) \leq \left(\bar{h}_{i,j}\right)(2V_0 + 2A_0) + \epsilon\text{''}. \tag{219}$$

Fixing that value for i, we now take $j \to \infty$,

$$\lim_{j\to\infty} d_{\mathcal{F}}(N', W_j) < \epsilon\text{''}. \tag{220}$$

We have the theorem as stated. □

Example 5.5. *In Figure 9 we see that a connected precompact exhaustion W_j of a standard flat two dimensional torus satisfying* $\mathrm{Vol}(W_j) \leq \mathrm{Vol}(N) \leq V_0$ *and*

$$\lim_{j\to\infty} \mathrm{Vol}(N \setminus W_j) = 0. \tag{221}$$

Observe that $(\bar{W}_j, d_{W_j})$ *need not have a Gromov-Hausdorff limit because balls of radius* $1/2$ *about the tips of the arms measured with respect to the intrinsic length metric* d_{W_j} *are disjoint and so the number of disjoint balls of radius* $1/2$ *is unbounded. According to the converse of Gromov Compactness Theorem, the number of disjoint balls in a sequence of compact metric spaces converging to a compact metric space is uniformly bounded above, so this sequence cannot converge* [Gro99].



FIGURE 9. W_1 with two arms is depicted in white on a black T^2. W_2 with four arms is depicted in light grey containing W_1. W_3 with eight arms is depicted in darker grey containing W_2 and W_1.

To find an example which also satisfies $\mathrm{Vol}(\partial W_j) \le A_0$, *we may construct a connected precompact exhaustion of a standard flat three dimensional torus where the arms are thin tubular neighborhoods of curves so that their lengths are still long enough to have disjoint balls but the areas of the boundaries of the arms are arbitrarily small.*

5.2. **Proof of Theorem 5.2.** In this subsection we prove Theorem 5.2. Keep in mind Remark 5.3. First we prove a short lemma which will be applied here and elsewhere:

Lemma 5.6. *Let* $M_i = (M, g_i)$ *be a sequence of Riemannian manifolds such that there is a closed subset,* S, *and a connected precompact exhaustion,* W_j, *of* $M \setminus S$ *satisfying (2) such that* g_i *converge smoothly to* g_∞ *on each* W_j. *If*

$$\mathrm{Vol}_{g_i}(M \setminus W_j) \le V_j \text{ where } \lim_{j\to\infty} V_j = 0 \tag{222}$$

then there exists a uniform $V_0 > 0$ *such that*

$$\mathrm{Vol}_{g_i}(M) < V_0. \tag{223}$$

Proof. Fix any W_j. Since g_i converges smoothly on W_j, $\mathrm{Vol}_{g_i}(W_j)$ must converge smoothly as well. So there exists $V_1 > 0$ such that $\mathrm{Vol}_{g_i}(W_j) \le V_1$. Thus we have

$$\mathrm{Vol}_{g_i}(M) = \mathrm{Vol}_{g_i}(W_j) + \mathrm{Vol}_{g_i}(M \setminus W_j) \le V_1 + V_j \tag{224}$$

and $\sup V_j < \infty$ because $\lim_{j\to\infty} V_j$ exists. □

We now prove Theorem 5.2:

Proof. By hypothesis (200) and Lemma 5.6 we have:

$$\mathrm{Vol}(M_i) \le V_0, \tag{225}$$

Next we prove that (W_j, g_∞) satisfy the hypothesis of Theorem 5.4. Observe that hypothesis (198) implies

$$\mathrm{Vol}_{g_\infty}(W_j) = \lim_{i\to\infty} \mathrm{Vol}_{g_i}(W_j) \le V_0, \tag{226}$$

while (199) implies

$$\mathrm{Vol}_{g_\infty}(\partial W_j) = \lim_{i\to\infty} \mathrm{Vol}_{g_i}(\partial W_j) \le A_0. \tag{227}$$

Finally

$$\begin{aligned}
\mathrm{diam}_N(N) &= \lim_{j\to\infty} \mathrm{diam}_N(W_j) \tag{228}\\
&\le \lim_{j\to\infty}\lim_{k\to\infty} \mathrm{diam}_{(W_k,g_\infty)}(W_j) \tag{229}\\
&\le \lim_{j\to\infty}\lim_{k\to\infty}\lim_{i\to\infty} \mathrm{diam}_{(W_k,g_i)}(W_j) \tag{230}\\
&\le \lim_{j\to\infty}\lim_{k\to\infty}\lim_{i\to\infty} \mathrm{diam}_{(M,g_i)}(W_j) + \lambda_{i,j,k} \tag{231}\\
&\le \limsup_{j\to\infty}\limsup_{k\to\infty}\limsup_{i\to\infty} D_0 + \lambda_{i,j,k} \tag{232}\\
&\le D_0 + \lambda_0. \tag{233}
\end{aligned}$$

Thus by Theorem 5.2 we have

$$d_{\mathcal{F}}((W_j, g_\infty), (N', d_\infty)) = F_j \text{ where } \lim_{j\to\infty} F_j = 0. \tag{234}$$

Next we will apply Theorem 4.6 to show $M_1 = (W_k, g_\infty)$ and $M_2 = (M, g_i)$ are close in the intrinsic flat sense by setting $U_1 = W_j \subset W_k$ and $U_2 = W_j \subset M$ for some well chosen $j < k$. Then the values in the hypothesis of the theorem are

$$\begin{aligned}
\epsilon &= \epsilon_{i,j} \text{ where } \lim_{i\to\infty} \epsilon_{i,j} = 0, && (235)\\
D_{U_2} &\le \operatorname{diam}_{(M,g_i)}(W_j) \le D_0 && (236)\\
D_{U_1} &\le \operatorname{diam}_{(W_k,g_i)}(W_k) \le D_0 + \lambda_0 && (237)\\
a &= a_{i,j,k} \le a_{i,j} = 2(D_0+\lambda_0)\arccos(1+\epsilon_{i,j})^{-1}/\pi && (238)\\
\lambda &= \lambda_{i,j,k} && (239)\\
h &= h_{i,j,k} \le \sqrt{\lambda_{i,j,k}(D_0+\lambda_0+\lambda_{i,j,k}/4)} && (240)\\
\bar{h} &= \bar{h}_{i,j,k} \le \max\{h_{i,j,k}, \sqrt{\epsilon_{i,j}^2 + 2\epsilon_{i,j}}(D_0+\lambda_0)\} && (241)
\end{aligned}$$

Thus

$$d_{\mathcal{F}}((W_k, g_\infty),(M,g_i)) \le \left(\bar{h}_{i,j,k} + a_{i,j,k}\right)\left(2V_0 + 2A_0\right) + 2V_j. \tag{242}$$

Combining this with (234) we have for any $j < k$,

$$d_{\mathcal{F}}((N, g_\infty),(M,g_i)) \le \left(\bar{h}_{i,j,k} + a_{i,j,k}\right)\left(2V_0 + 2A_0\right) + 2V_j + F_k. \tag{243}$$

Taking the limsup as $k \to \infty$ we have

$$\begin{aligned}
d_{\mathcal{F}}((N,g_\infty),(M,g_i)) &\le \limsup_{k\to\infty}\left(\bar{h}_{i,j,k} + a_{i,j,k}\right)\left(2V_0+2A_0\right) + 2V_j && (244)\\
&\le \left(\bar{h}_{i,j} + a_{i,j}\right)\left(2V_0+2A_0\right) + 2V_j, && (245)
\end{aligned}$$

where

$$\begin{aligned}
\bar{h}_{i,j} &= \limsup_{k\to\infty} \bar{h}_{i,j,k} && (246)\\
&\le \max\{h_{i,j}, \sqrt{\epsilon_{i,j}^2 + 2\epsilon_{i,j}}(D_0+\lambda_0)\}, && (247)
\end{aligned}$$

and

$$h_{i,j} = \limsup_{k\to\infty} h_{i,j,k} \le \sqrt{\lambda_{i,j}(D_0+\lambda_0+\lambda_{i,j}/4)}. \tag{248}$$

Recall that for any fixed j, $\lim_{i\to\infty}\epsilon_{i,j} = 0$, thus $\lim_{i\to\infty} a_{i,j} = 0$ as well. By the hypothesis $\limsup_{i\to\infty}\lambda_{i,j} = \lambda_j$ so

$$\bar{h}_j = \limsup_{i\to\infty} \bar{h}_{i,j} \le \sqrt{\lambda_j(D_0+\lambda_0+\lambda_j/4)}. \tag{249}$$

Thus

$$\limsup_{i\to\infty} d_{\mathcal{F}}((N,g_\infty),(M,g_i)) \le \bar{h}_j(2V_0+2A_0) + 2V_j \qquad \forall j \in \mathbb{N}. \tag{250}$$

By the hypothesis, taking $j \to \infty$ we have,

$$\limsup_{i\to\infty} d_{\mathcal{F}}((N,g_\infty),(M,g_i)) = 0 \tag{251}$$

□

5.3. **Codimension 2 Singular Sets.** The following lemma combined with Theorem 5.2 completes the proof of Theorem 1.3.

Lemma 5.7. *M compact and* $\operatorname{diam}_{g_\infty}(M \setminus S) < \infty$ *and if S is a submanifold of codimension* 2 *then any connected precompact exhaustion, W_j, of $M \setminus S$ is uniformly well embedded.*

Proof. I claim for fixed i, j,

$$\lambda_{i,j} = \limsup_{k\to\infty} \lambda_{i,j,k} = 0. \tag{252}$$

Suppose not.

Let $x_{i,j,k}, y_{i,j,k} \subset \bar{W}_j$ achieve to supremum in the definition of $\lambda_{i,j}$.

Since $\bar{W}_j$ is compact, a subsequence as $k \to \infty$ converges to $x_{i,j}, y_{i,j} \subset \bar{W}_j$. Let $\gamma_{i,j}$ be a minimizing geodesic between these points in M with respect to g_i. Since S is a submanifold of codimension 2, we can find a curve $C_{i,j} : [0,1] \to M \setminus S$ between these points such that

$$L_{g_i}(C_{i,j}) \le d_{M,g_i}(x_{i,j}, y_{i,j}) + \lambda_{i,j}/5 \tag{253}$$

by sliding $\gamma_{i,j}$ over slightly to avoid S.

Let k be chosen from the subsequence sufficiently large that

$$C_{i,j}([0,1]) \subset U_k d_{(\bar{U}_j,g_i)}(x_{i,j,k}, x_{i,j}) < \lambda_{i,j}/10 d_{(\bar{U}_j,g_i)}(y_{i,j,k}, y_{i,j}) < \lambda_{i,j}/10 \tag{254}$$

Thus

$$\begin{aligned}
d_{(\bar{U}_k,g_i)}(x_{i,j,k}, y_{i,j,k}) &\le d_{(\bar{U}_k,g_i)}(x_{i,j,k}, x_{i,j}) + d_{(\bar{U}_k,g_i)}(x_{i,j}, y_{i,j}) + d_{(\bar{U}_k,g_i)}(y_{i,j}, y_{i,j,k}) \\
&\le d_{(\bar{U}_j,g_i)}(x_{i,j,k}, x_{i,j}) + L(C_{i,j}) + d_{(\bar{U}_j,g_i)}(y_{i,j}, y_{i,j,k}) \\
&\le \lambda_{i,j}/10 + d_{M,g_i}(x_{i,j}, y_{i,j}) + \lambda_{i,j}/5 + \lambda_{i,j}/10 \\
&\le 2\lambda_{i,j}/5 + d_{M,g_i}(x_{i,j}, x_{i,j,k}) + d_{M,g_i}(x_{i,j,k}, y_{i,j,k}) + d_{M,g_i}(y_{i,j,k}, y_{i,j}) \\
&\le 2\lambda_{i,j}/5 + d_{U_j,g_i}(x_{i,j}, x_{i,j,k}) + d_{M,g_i}(x_{i,j,k}, y_{i,j,k}) + d_{U_j,g_i}(y_{i,j,k}, y_{i,j}) \\
&\le 3\lambda_{i,j}/5 + d_{M,g_i}(x_{i,j,k}, y_{i,j,k}) \\
&\le 3\lambda_{i,j}/5 + d_{U_k,g_i}(x_{i,j,k}, y_{i,j,k}) - \lambda_{i,j,k}
\end{aligned}$$

by the choice of $x_{i,j,k}$ and $y_{i,j,k}$. This is a contradiction.

Next we must show

$$\limsup_{j\to\infty} \limsup_{k\to\infty} \limsup_{i\to\infty} \lambda_{i,j,k} \le \lambda_0. \tag{255}$$

Observe that

$$\lambda_{i,j,k} \le \bar{\lambda}_{i,j,k} = \operatorname{diam}_{(W_k,g_i)}(W_j) \tag{256}$$

Since $g_i \to g_\infty$ on W_k we know

$$\limsup_{i\to\infty} \lambda_{i,j,k} \le \mathrm{diam}_{(W_k,g_\infty)}(W_j). \tag{257}$$

We see by the exhaustion that

$$\lim_{k\to\infty} \limsup_{i\to\infty} \lambda_{i,j,k} \le \mathrm{diam}_{(M\setminus S,g_\infty)}(W_j) \tag{258}$$

and so

$$\limsup_{j\to\infty} \limsup_{k\to\infty} \limsup_{i\to\infty} \lambda_{i,j,k} \le \mathrm{diam}_{g_\infty}(M \setminus S). \tag{259}$$

□

6. INTRINSIC FLAT TO GH CONVERGENCE

There are occasions where one has volume controls as in Theorem 5.2 but one would like to obtain a Gromov-Hausdorff limit. That is not always possible. Example 3.10 has no Gromov-Hausdorff limit despite satisfying the conditions of Theorem 5.2. In Example 3.9 the Gromov-Hausdorff limits and intrinsic flat limits do not agree. However the first author and Stefan Wenger have shown in [SW11] that the Gromov-Hausdorff and intrinsic flat limits agree when the sequence of manifolds has nonnegative Ricci curvature or a uniform contractibility function:

Definition 6.1. *A function $\rho : [0, r_0] \to [0, \infty)$ is a contractibility function for a manifold M with metric g if every ball $B_p(r)$ is contractible within $B_p(\rho(r))$.*

We review these results in the first subsection.

In the second subsection, we apply the results in [SW11] on sequences of manifolds with a uniform contractibility function, proving Theorem 6.7 and Theorem 6.6.

In the third subsection we use additional properties of of manifolds with Ricci curvature bounds to prove additional theorems about Gromov-Hausdorff limits inspired by the techniques in [SW11]. In particular we prove Theorem 6.10 and Theorem 1.2.

6.1. Review of Convergence Theorems.

First recall that Gromov proved a sequence of Riemannian manifolds has a subsequence converging in the Gromov-Hausdorff sense if there is a uniform bound on the number of disjoint balls of radius r that fit in the space [Gro99]. This lead to two compactness theorems:

Theorem 6.2 (Gromov)**.** [Gro99] *A sequence of compact Riemannian manifolds, (M_j, g_j), such that* $\mathrm{diam}(M_j) \le D$ *and $Ricci_{M_j} \ge -H$, has a subsequence converging in the Gromov-Hausdorff sense to a metric space (X, d).*

Theorem 6.3 (Greene-Petersen)**.** [GPV92] *A sequence of compact Riemannian manifolds,* (M_j, g_j)*, such that* $\mathrm{Vol}(M_j) \le V$ *and such that there is a uniform contractibility function,* $\rho : [0, r_0] \to [0, \infty)$*, for all the* M_j*, has a subsequence converging in the Gromov-Hausdorff sense to a metric space* (X, d)*.*

See Definition 6.1.

In [SW10] the following theorems were proven which can be applied to deduce information about the Gromov-Hausdorff limit of a sequence.

Theorem 6.4 (Sormani-Wenger)**.** *If a sequence of oriented compact Riemannian manifolds,* (M_j, g_j)*, such that* $\mathrm{diam}(M_j) \le D$ *and* $Ricci_{M_j} \ge 0$ *and* $vol(M_j) \ge V_0$ *converges in the Gromov-Hausdorff sense to* (X, d)*, then it converges in the intrinsic flat Sense to* (X, d, T) *(c.f. Theorem 4.16 of* [SW11]*).*

This theorem is conjectured to hold with uniform lower bounds on Ricci curvature [SW11].

Theorem 6.5 (Sormani-Wenger)**.** *If a sequence of oriented compact Riemannian manifolds,* (M_j, g_j)*, with a uniform linear contractibility function,* $\rho : [0, \infty) \to [0, \infty)$ *and a uniform upper bound on volume,* $\mathrm{Vol}(M_j) \le V$*, converges in the Gromov-Hausdorff sense to* (X, d)*, then it converges in the intrinsic flat Sense to* (X, d, T) *(c.f. Theorem 4.14 of* [SW11]*).*

Recall that, in general, the intrinsic flat limits and Gromov-Hausdorff limits need not agree [Examples 3.3 and 3.9] because intrinsic flat limits do not include points with 0 density as in (26). In fact intrinsic flat limits may exist when Gromov-Hausdorff limits do not [Example 3.10].

6.2. **Sequences with Uniform Contractibility Functions.**

Recall Definition 6.1. Here we apply the results in [SW11] on sequences of manifolds with a uniform contractibility function, stating and proving Theorem 6.7 and Theorem 6.6. Recall Definitions 1.1 and 5.1.

Theorem 6.6. *Let* $M_i = (M, g_i)$ *be a sequence of Riemannian manifolds with a uniform linear contractibility function,* ρ*, which converges smoothly away from a singular set,* S*, of codimension two. If there is a connected precompact exhaustion of* $M \setminus S$ *as in (2) satisfying the volume conditions*

$$\mathrm{Vol}_{g_i}(\partial W_j) \le A_0 \tag{260}$$

and

$$\mathrm{Vol}_{g_i}(M \setminus W_j) \le V_j \text{ where } \lim_{j\to\infty} V_j = 0. \tag{261}$$

then

$$\lim_{j\to\infty} d_{GH}(M_j, N) = 0 \tag{262}$$

where N *is the settled and metric completion of* $(M \setminus S, g_\infty)$*.*

Theorem 6.7. *Let $M_i = (M, g_i)$ be a sequence of Riemannian manifolds with a uniform linear contractibility function, ρ, which converges smoothly away from a singular set, S. If there is a uniformly well embedded connected precompact exhaustion of $M \setminus S$ as in (2) satisfying the volume conditions (260) and (261) then*

$$\lim_{j\to\infty} d_{GH}(M_j, N) = 0 \tag{263}$$

where N is the settled and metric completion of $(M \setminus S, g_\infty)$.

Remark 6.8. *Example 3.9 has no uniform linear contractibility near the singular set and the Gromov-Hausdorff limit does not agree with the intrinsic flat limit. Examples 3.10 and 3.11, also satisfy all the conditions of Theorem 6.6 and 6.7 except the existence of a uniform linear contractibility function. They have no Gromov-Hausdorff limit.*

The excess volume bound in (4) is shown to be necessary in Example 3.7 and Example 3.8. The codimension two condition of Theorem 6.6 and the uniform embeddedness hypothesis of Theorem 6.7 are seen to be necessary in Example 3.15. We believe we have an example proving the necessity of the uniform bound on the boundary volumes, (3), and discuss this in Remark 3.13.

Remark 6.9. *It would be interesting to see whether the requirement that the contractibility function is linear is a necessary condition. One might consider adapting the Example by Schul and Wenger in the appendix of* [SW10] *to prove this.*

Proof. By Lemma 5.6, we have

$$\mathrm{Vol}(M_i) \le V_0. \tag{264}$$

This combined with the uniform contractibility function allows us to apply the Greene-Petersen Compactness Theorem. In particular we have a uniform upper bound on diameter:

$$\mathrm{diam}(M_i) \le D_0, \tag{265}$$

We may now apply Theorem 5.2 to obtain

$$\lim_{j\to\infty} d_{\mathcal{F}}(M_j, N') = 0 \tag{266}$$

We then apply Theorem 6.5 to see that the flat limit and Gromov-Hausdorff limits agree due to the existence of the uniform linear contractibility function and the fact that the volume is bounded below uniformly by the smooth limit. In particular the metric completion and the settled completion agree. □

We now easily prove Theorem 6.6:

Proof. This theorem follows from Theorem 6.7 combined with Lemma 5.7. □

6.3. **Ricci curvature bounded below.** In this subsection we use additional properties of of manifolds with Ricci curvature bounds to prove additional theorems about Gromov-Hausdorff limits inspired by the techniques in [SW11]. In particular we prove Theorem 6.10 and Theorem 1.2. Recall Definitions 1.1, 2 and 5.1.

Theorem 6.10. *Let $M_i = (M, g_i)$ be a sequence of Riemannian manifolds with uniform lower Ricci curvature bounds,*

$$\mathrm{Ricci}_{\mathrm{g_i}}(\mathrm{V},\mathrm{V}) \geq (\mathrm{n}-1)\mathrm{H}\,\mathrm{g_i}(\mathrm{V},\mathrm{V}) \qquad \forall \mathrm{V} \in \mathrm{TM_i} \tag{267}$$

which converges smoothly away from a singular set, $S \subset M$. If there is a uniformly well embedded connected precompact exhaustion of $M \setminus S$ as in (2) satisfying the volume conditions, (3) and (4) then

$$\lim_{j\to\infty} d_{GH}(M_j, N) = 0 \tag{268}$$

where N is the settled and metric completion of $(M \setminus S, g_\infty)$.

When $H = 0$ this theorem is an immediate consequence of Theorem 5.2 assuming in addition that

$$\mathrm{diam}_{M_i}(W_j) \leq D_0 \qquad \forall i \geq j, \tag{269}$$

To remove these unnecessary conditions we prove the following lemma and proposition

Lemma 6.11. *Suppose we have a sequence of manifolds, $M_j = (M, g_j)$ with a uniform lower bound on Ricci curvature and*

$$\mathrm{Vol}(M_j) \leq V_0 \tag{270}$$

converging smoothly away from a singular set to $(M \setminus S, g_\infty)$ then

$$\mathrm{diam}_{M_i}(W_j) \leq \mathrm{diam}(M_i) \leq D_0 \qquad \forall i \geq j, \tag{271}$$

Proof. Suppose not. Let $p \in W \subset \bar{W} \subset M \setminus S$ where W is precompact and let $q_j \in M_j$ such that $d_j = d_j(p, q_j) \to \infty$. By smooth convergence on W, there exists $r_0 > 0$ such that $B_p(r_0) \subset M_j$ smoothly converge to a ball in a smooth Riemannian manifold. In particular $\mathrm{Vol}_{g_j}(B_p(r_0)) \geq V_1$. Then

$$\begin{aligned}
V_0 \;&\geq\; \mathrm{Vol}(B_q(d_j - r_0)) &(272)\\
&\geq\; \frac{(d_j - r_0)^m}{d_j^m - (d_j - r_0)^m}\,\mathrm{Vol}(Ann_q(d_j - r_0, d_j)) &(273)\\
&\geq\; \frac{(d_j - r_0)^m}{d_j^m - (d_j - r_0)^m}\,\mathrm{Vol}(B_p(r_0)) &(274)\\
&\geq\; \frac{(d_j - r_0)^m}{2m d_j^{m-1} r_0} V_1 &(275)
\end{aligned}$$

which gives a contradiction as $d_j \to \infty$. □

This proposition extending the results in [SW10]:

Proposition 6.12. *Suppose we have a sequence of manifolds, $M_j = (M, g_j)$ with a uniform lower bound on Ricci curvature and*

$$\mathrm{Vol}(M_j) \le V_0 \tag{276}$$

converging smoothly away from a singular set to $(M \setminus S, g_\infty)$. Suppose also that (M, g_j) converge in the intrinsic flat sense to N' where N' is the settled completion of $(M \setminus S, g_\infty)$. Then

$$d_{GH}(\bar{M}_j, \bar{N}) \to 0 \tag{277}$$

and $\bar{N} = N'$.

Proof. By Lemma 6.11, $\mathrm{diam}(M_j) \le D_0$. By Gromov's Compactness theorem, we know that a subsequence of the Riemannian manifolds $M_j = (M, g_j)$ converge to a compact metric space (Y, d). Thus a subsequence of the manifolds converges in the intrinsic flat sense to an integral current space, (X, d, T), where $X \subset Y$.

By Theorem 5.2 and the fact that intrinsic flat limits are unique we know that the settled completion of $(M \setminus S, g_\infty)$ is (X, d, T). In particular one needs no subsequence to obtain the flat limit.

In the case where the sequence of metrics has nonnegative Ricci curvature, Theorem 6.4 implies that $X = Y$. In particular the settled completion is the metric completion and so the Gromov-Hausdorff limit is the metric completion of $(M \setminus S, g_\infty)$ and no subsequence was needed.

When the sequence of manifolds has a negative uniform lower bound on Ricci curvature, we may imitate the proof of Theorem 6.4. We must show that every $y \in Y$ lies in the settled completion of $(M \setminus S, g_\infty)$.

First observe that by the smooth convergence of g_j away from S, we know the volumes are uniformly bounded below:

$$\mathrm{Vol}_j(M_j) \ge V_0. \tag{278}$$

Thus we can apply the noncollapsing theory of Cheeger-Colding [CC97] to see that after possibly taking another subsequence of (M, g_j) we can control the volumes of the limit spaces balls: For all $y \in Y$, there exists $y_j \in M$ such that

$$\lim_{j \to \infty} \mathrm{Vol}_j(B_{y_j}(r)) = \mathcal{H}_m(B_y(r)) \ge V_0(r/D)^m > 0, \tag{279}$$

where $B_y(r) \subset Y$ and $\mathcal{H}_m$ is the m dimensional volume. In particular, for j sufficiently large

$$\mathrm{Vol}_j(B_{y_j}(r)) \ge (V_0/2)(r/D)^m > 0. \tag{280}$$

Now we choose i sufficiently large (depending on r), so that

$$\epsilon_i < (V_0/4)(r/D)^m. \tag{281}$$

Then

$$\mathrm{Vol}_j(U_i \cap B_{y_j}(r)) > (V_0/4)(r/D)^m > 0. \tag{282}$$

Thus there exists

$$z_{r,j} \in U_i \cap B_{y_j}(r) \subset M \setminus S. \tag{283}$$

and

$$\mathrm{Vol}_j(U_i \cap B_{z_{r,j}}(2r)) \geq \mathrm{Vol}_j(U_i \cap B_{y_j}(r)). \tag{284}$$

Since $z_{r,j} \subset U_i \subset \bar{U}_i$, a subsequence of the $z_{r,j}$ converge to $z_{r,\infty} \in \bar{U}_i \subset U_{i+1}$. Since g_j converge smoothly to g_∞ on U_{i+1},

$$\mathrm{Vol}_j(U_i \cap B_{z_{r,j}}(2r)) \to \mathrm{Vol}_\infty(U_i \cap B_{z_{r,\infty}}(2r)). \tag{285}$$

Thus

$$\mathrm{Vol}_\infty(U_i \cap B_{z_{r,\infty}}(2r)) \geq (V_0/4)(r/D)^m > 0 \tag{286}$$

Note that by (283), taking the Gromov-Hausdorff limit we see that

$$d(z_{r,\infty}, y) < r. \tag{287}$$

So y is in the metric completion of $(M \setminus S, g_\infty)$. Furthermore

$$\begin{aligned} \mathrm{Vol}(B_y(3r) \cap (M \setminus S)) &\geq \mathrm{Vol}(B_y(3r) \cap U_i) && (288)\\ &\geq \mathrm{Vol}(B_{z_{r,\infty}}(2r) \cap U_i) && (289)\\ &\geq (V_0/4)(r/D)^m > 0 && (290)\end{aligned}$$

so y is in the settled completion of $(M \setminus S, g_\infty)$.

In particular the settled completion is the metric completion and so the Gromov-Hausdorff limit is the metric completion of $(M \setminus S, g_\infty)$ and no subsequence was needed. □

We may now prove Theorem 6.10:

Proof. By Lemma 6.11, $\mathrm{diam}(M_j) \leq D_0$. This combined with the hypothesis (3) and (4), allows us to apply Theorem 5.2. So (M_i, g_i) has an intrinsic flat limit and that this intrinsic flat limit is the settled completion of $(M \setminus S, g_\infty)$. By Lemma 5.6, we have (276). Thus by Proposition 6.12, the Gromov-Hausdorff and Intrinsic Flat limits agree. □

We now prove Theorem 1.2 which was stated in the introduction:

Proof. This theorem follows from Theorem 6.10 combined with Lemma 5.7. □

Remark 6.13. *The Ricci curvature condition is necessary in Theorem 1.2 as can be seen in Example 3.9 and in Examples 3.10 and 3.11, which have no Gromov-Hausdorff limit. The excess volume bound in (4) is shown to be necessary in Example 3.7 and Example 3.8, which has no Gromov-Hausdorff limit. All these examples satisfy the uniform embeddedness hypothesis of Theorem 6.10 and demonstrate the* *necessity of these conditions in that theorem as well. We were unable to find an example proving the necessity of the uniform bound on the boundary volumes, (3), but discuss this in Remark 3.14. The codimension two condition of Theorem 1.2 and the uniform embeddedness hypothesis of Theorem 6.10 are seen to be necessary for their respective theorems in Example 3.15.*

REFERENCES

[AK00] Luigi Ambrosio and Bernd Kirchheim, *Currents in metric spaces*, Acta Math. **185** (2000), no. 1, 1–80. MR MR1794185 (2001k:49095)

[And89] Michael T. Anderson, *Ricci curvature bounds and Einstein metrics on compact manifolds*, J. Amer. Math. Soc. **2** (1989), no. 3, 455–490. MR 999661 (90g:53052)

[Ban90] Shigetoshi Bando, *Bubbling out of Einstein manifolds*, Tohoku Math. J. (2) **42** (1990), no. 2, 205–216. MR 1053949 (92a:53065a)

[Bas] Jorge Basilio, *The scalable intrinsic flat distance*, preprint in progress.

[BKN89] Shigetoshi Bando, Atsushi Kasue, and Hiraku Nakajima, *On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth*, Invent. Math. **97** (1989), no. 2, 313–349. MR 1001844 (90c:53098)

[CC97] Jeff Cheeger and Tobias H. Colding, *On the structure of spaces with Ricci curvature bounded below. I*, J. Differential Geom. **46** (1997), no. 3, 406–480. MR MR1484888 (98k:53044)

[EGZ09] Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi, *Singular Kähler-Einstein metrics*, J. Amer. Math. Soc. **22** (2009), no. 3, 607–639. MR 2505296 (2010k:32031)

[FF60] Herbert Federer and Wendell H. Fleming, *Normal and integral currents*, Ann. of Math. (2) **72** (1960), 458–520. MR MR0123260 (23 #A588)

[GPV92] Robert E. Greene and Peter Petersen V, *Little topology, big volume*, Duke Math. J. **67** (1992), no. 2, 273–290.

[Gro83] Mikhael Gromov, *Filling Riemannian manifolds*, J. Differential Geom. **18** (1983), no. 1, 1–147.

[Gro99] Misha Gromov, *Metric structures for Riemannian and non-Riemannian spaces*, Progress in Mathematics, vol. 152, Birkhäuser Boston Inc., Boston, MA, 1999, Based on the 1981 French original [MR0682063 (85e:53051)], With appendices by M. Katz, P. Pansu and S. Semmes, Translated from the French by Sean Michael Bates. MR MR1699320 (2000d:53065)

[Hua09] Hong Huang, *Convergence of Einstein 4-orbifolds*, Acta Math. Sinica (Chin. Ser.) **52** (2009), no. 1, 205–208. MR 2513581 (2010a:53068)

[Lak] Sajjad Lakzian, *Doctoral dissertation*, In progress.

[LS11] Dan A. Lee and Christina Sormani, *Stability of the positive mass theorem for rotationally symmetric riemannian manifolds*, preprint on arxiv (2011).

[LSar] ______, *Almost inequality in the Penrose equality for rotationally symmetric manifolds*, Annales Henri Poincare (to appear).

[RZ] Ruan and Zhang, *Convergence of calabi-yau manifolds*, preprint on arxiv.

[Ses04] Natasa Sesum, *Convergence of Kähler-Einstein orbifolds*, J. Geom. Anal. **14** (2004), no. 1, 171–184. MR 2030580 (2004i:32040)

[SW10] Christina Sormani and Stefan Wenger, *Weak convergence and cancellation, appendix by Raanan Schul and Stefan Wenger*, Calculus of Variations and Partial Differential Equations **38** (2010), no. 1-2.

[SW11] ______, *Intrinsic flat convergence of manifolds and other integral current spaces*, Journal of Differential Geometry **87** (2011).

[Tia90] G. Tian, *On Calabi's conjecture for complex surfaces with positive first Chern class*, Invent. Math. **101** (1990), no. 1, 101–172. MR 1055713 (91d:32042)

[Tos09] Valentino Tosatti, *Limits of Calabi-Yau metrics when the Kähler class degenerates*, J. Eur. Math. Soc. (JEMS) **11** (2009), no. 4, 755–776. MR 2538503 (2010j:32039)

CUNY Graduate Center and Lehman College
E-mail address: `SLakzian@gc.cuny.edu`

CUNY Graduate Center and Lehman College
E-mail address: `sormanic@member.ams.org`